

On general multiple Eulerian integrals and multivariable Aleph-function

F.Y. AYANT¹

¹ Teacher in High School , France
 E-mail : fredericayant@gmail.com

ABSTRACT

The object of this paper is first, to evaluate a general multiple Eulerian integrals with general integrands involving the product of a multivariable Aleph-function and generalized hypergeometric function with general arguments. The second multiple Eulerian integral contain a general class of polynomials the general polynomials set and multivariable Aleph-function with general arguments. Our integral formulas are interesting and unified nature. We will study the case of multivariable I-function defined by Sharma et al [7].

Keywords : Multivariable Aleph-functions, multivariable I-function, class of polynomial, general polynomials set, multiple Eulerian integral

2010 Mathematics Subject Classification. 33C99, 33C60, 44A20

1. Introduction

In this paper, first, to evaluate a general multiple Eulerian integrals with general integrands involving the product of a multivariable Aleph-function and generalized hypergeometric function with general arguments. The second multiple Eulerian integral contain a general class of polynomials the general polynomial set and multivariable Aleph-function with general arguments. Our integral formulas are interesting and unified nature. We will study the case of multivariable I-function defined by Sharma et al [7].

The Aleph-function of several variables generalize the multivariable I-function defined by Sharma and Ahmad [7] , itself is an a generalisation of G and H-functions of several variables. The multiple Mellin-Barnes integral occurring in this paper will be referred to as the multivariables Aleph-function throughout our present study and will be defined and represented as follows.

$$\text{We have : } \aleph(z_1, \dots, z_r) = \aleph_{p_i, q_i, \tau_i; R: p_i(1), q_i(1), \tau_i(1); R^{(1)}; \dots; p_i(r), q_i(r), \tau_i(r); R^{(r)}}^{0, n: m_1, n_1, \dots, m_r, n_r} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \right)$$

$$[(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, n_1}, [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n_1+1, p_i}]:$$

$$\dots, [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m_1+1, q_i}]:$$

$$\left(\begin{matrix} [(c_j^{(1)}, \gamma_j^{(1)})_{1, n_1}, [\tau_{i(1)}(c_{ji}^{(1)}, \gamma_{ji}^{(1)})_{n_1+1, p_i^{(1)}}]; \dots; [(c_j^{(r)}, \gamma_j^{(r)})_{1, n_r}, [\tau_{i(r)}(c_{ji}^{(r)}, \gamma_{ji}^{(r)})_{n_r+1, p_i^{(r)}}] \\ [(d_j^{(1)}, \delta_j^{(1)})_{1, m_1}, [\tau_{i(1)}(d_{ji}^{(1)}, \delta_{ji}^{(1)})_{m_1+1, q_i^{(1)}}]; \dots; [(d_j^{(r)}, \delta_j^{(r)})_{1, m_r}, [\tau_{i(r)}(d_{ji}^{(r)}, \delta_{ji}^{(r)})_{m_r+1, q_i^{(r)}}] \end{matrix} \right)$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} ds_1 \dots ds_r \tag{1.1}$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^n \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} s_k)]} \tag{1.2}$$

$$\text{and } \theta_k(s_k) = \frac{\prod_{j=1}^{m_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{n_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i(k)=1}^{R^{(k)}} [\tau_{i(k)} \prod_{j=m_k+1}^{q_i(k)} \Gamma(1 - d_{ji}^{(k)} + \delta_{ji}^{(k)} s_k) \prod_{j=n_k+1}^{p_i(k)} \Gamma(c_{ji}^{(k)} - \gamma_{ji}^{(k)} s_k)]} \tag{1.3}$$

Suppose, as usual, that the parameters

$$a_j, j = 1, \dots, p; b_j, j = 1, \dots, q;$$

$$c_j^{(k)}, j = 1, \dots, n_k; c_{j i^{(k)}}^{(k)}, j = n_k + 1, \dots, p_{i^{(k)}};$$

$$d_j^{(k)}, j = 1, \dots, m_k; d_{j i^{(k)}}^{(k)}, j = m_k + 1, \dots, q_{i^{(k)}};$$

$$\text{with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$$

are complex numbers, and the α 's, β 's, γ 's and δ 's are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{j i}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{j i^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{j i}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{j i^{(k)}}^{(k)} \leq 0 \quad (1.4)$$

The real numbers τ_i are positives for $i = 1$ to R , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary, ensure that the poles of $\Gamma(d_j^{(k)} - \delta_j^{(k)} s_k)$ with $j = 1$ to m_k are separated from those of $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$ with $j = 1$ to n and $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$ with $j = 1$ to n_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{j i}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{j i}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{j i^{(k)}}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{j i^{(k)}}^{(k)} > 0, \text{ with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \quad (1.5)$$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the asymptotic expansion in the following convenient form :

$$\aleph(y_1, \dots, y_r) = O(|y_1|^{\alpha_1}, \dots, |y_r|^{\alpha_r}), \max(|y_1|, \dots, |y_r|) \rightarrow 0$$

$$\aleph(y_1, \dots, y_r) = O(|y_1|^{\beta_1}, \dots, |y_r|^{\beta_r}), \min(|y_1|, \dots, |y_r|) \rightarrow \infty$$

where $k = 1, \dots, r : \alpha_k = \min[Re(d_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, m_k$ and

$$\beta_k = \max[Re((c_j^{(k)} - 1) / \gamma_j^{(k)})], j = 1, \dots, n_k$$

We will use these following notations in this paper

$$U = p_i, q_i, \tau_i; R ; V = m_1, n_1; \dots ; m_r, n_r \quad (1.6)$$

$$W = p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}, \dots, p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)} \quad (1.7)$$

$$A = \{(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1,n}\}, \{\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1,p_i}\} \quad (1.8)$$

$$B = \{\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1,q_i}\} \quad (1.9)$$

$$C = \{(c_j^{(1)}; \gamma_j^{(1)})_{1,n_1}\}, \tau_{i(1)}(c_{ji(1)}^{(1)}; \gamma_{ji(1)}^{(1)})_{n_1+1,p_{i(1)}}, \dots, \{(c_j^{(r)}; \gamma_j^{(r)})_{1,n_r}\}, \tau_{i(r)}(c_{ji(r)}^{(r)}; \gamma_{ji(r)}^{(r)})_{n_r+1,p_{i(r)}} \quad (1.10)$$

$$D = \{(d_j^{(1)}; \delta_j^{(1)})_{1,m_1}\}, \tau_{i(1)}(d_{ji(1)}^{(1)}; \delta_{ji(1)}^{(1)})_{m_1+1,q_{i(1)}}, \dots, \{(d_j^{(r)}; \delta_j^{(r)})_{1,m_r}\}, \tau_{i(r)}(d_{ji(r)}^{(r)}; \delta_{ji(r)}^{(r)})_{m_r+1,q_{i(r)}} \quad (1.11)$$

The multivariable Aleph-function write :

$$\aleph(z_1, \dots, z_r) = \aleph_{U:W}^{0,n:V} \left(\begin{array}{c|c} z_1 & \text{A : C} \\ \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ z_r & \text{B : D} \end{array} \right) \quad (1.12)$$

Srivastava [8] introduced the general class of polynomials :

$$S_N^M(x) = \sum_{K=0}^{[N/M]} \frac{(-N)_{MK}}{K!} A_{N,k} x^K, N = 0, 1, 2, \dots \quad (1.13)$$

where M is an arbitrary positive integer and the coefficient $A_{N,k}$ are arbitrary constants, real or complex.

By suitably specialized the coefficient $A_{N,k}$ the polynomials $S_N^M(x)$ can be reduced to the classical orthogonal polynomials such as Jacobi, Hermite, Legendre and Laguerre polynomials etc.

2. Sequence of function

Agarwal and Chaubey [1], Salim [6] and several others have studied a general sequence of functions. In the present document we shall study the following useful series formula for a general sequence of functions.

$$R_n^{\alpha,\beta}[x; E, F, g, h; p, q; \gamma; \delta; e^{-sx^r}] = \sum_{w,v,u,t,e,k_1,k_2} \psi(w, v, u, t, e, k_1, k_2) x^R \quad (2.1)$$

$$\text{where } \sum_{w,v,u,t,e,k_1,k_2} = \sum_{w=0}^{\infty} \sum_{v=0}^n \sum_{u=0}^v \sum_{t=0}^n \sum_{c=0}^t \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \quad (2.2)$$

and the infinite series on the right side (2.1) is absolutely convergent, $R = ln + qv + pt + rw + k_1r + k_2q$ (2.3)

$$\text{and } \psi(w, v, u, t, e, k_1, k_2) = \frac{(-)^{t+w+k_2} (-v)_u (-t)_e (\alpha)_t l^n}{w!v!u!t!e!K_n k_1!k_2!} \frac{s^{w+k_1} F^{\gamma n-t}}{(1-\alpha-t)_e} (\alpha-\gamma n)_e$$

$$(-\beta-\delta n)_v g^{v+k_2} h^{\delta n-v-k_2} (v-\delta n)_{k_2} E^t \left(\frac{pe + rw + \lambda + qn}{l} \right)_n \quad (2.4)$$

where K_n is a sequence of constants.

By suitably specializing the parameters involving in (2.1), a general sequence of function reduced to generalized polynomials set studied by Raizada [5], a class of polynomials introduced by Fujiwara [2] and several others authors.

3. Integral representation of generalized hypergeometric function

The following generalized hypergeometric function in terms of multiple contour integrals is also required [10 ,page 39 eq .30]

$$\frac{\prod_{j=1}^P \Gamma(A_j)}{\prod_{j=1}^Q \Gamma(B_j)} {}_P F_Q [(A_P); (B_Q); -(x_1 + \dots + x_r)]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \frac{\prod_{j=1}^P \Gamma(A_j + s_1 + \dots + s_r)}{\prod_{j=1}^Q \Gamma(B_j + s_1 + \dots + s_r)} \Gamma(-s_1) \dots \Gamma(-s_r) x_1^{s_1} \dots x_r^{s_r} ds_1 \dots ds_r \quad (3.1)$$

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of $\Gamma(A_j + s_1 + \dots + s_r)$ are separated from those of $\Gamma(-s_j), j = 1, \dots, r$. The above result (1.23) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of $\Gamma(-s_j), j = 1, \dots, r$

The equivalent form of Eulerian beta integral is :

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} dt = (b-a)^{\alpha+\beta} B(\alpha, \beta) (Re(\alpha) > 0, Re(\beta) > 0, a \neq b) \quad (3.2)$$

4. First integral

We note :

$$V_1 = V; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0; W_1 = W; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1 \quad (4.1)$$

$$C_1 = C; (1, 0), \dots, (1, 0); (1, 0), \dots, (1, 0); D_1 = D; (0, 1), \dots, (0, 1); (0, 1), \dots, (0, 1) \quad (4.2)$$

$$A^* = [1 + \sigma'_i; \rho_i^{(1,1)}, \dots, \rho_i^{(1,r)}, \tau_i^{(1,1)}, \dots, \tau_i^{(1,l)}, 1, 0, \dots, 0]_{1,s}, \dots,$$

$$[1 + \sigma_i^{(T)}; \rho_i^{(T,1)}, \dots, \rho_i^{(T,r)}, \tau_i^{(T,1)}, \dots, \tau_i^{(T,l)}, 1, 0, \dots, 0]_{1,s}, [1 - A_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0]_{1,P},$$

$$[1 - \alpha_i; \delta'_i, \dots, \delta_i^{(r)}, \mu'_i, \dots, \mu_i^{(l)}, 1, \dots, 1, 0, \dots, 0]_{1,s},$$

$$[1 - \beta_i; \eta'_i, \dots, \eta_i^{(r)}, \theta'_i, \dots, \theta_i^{(l)}, 0, \dots, 0, 1, \dots, 1]_{1,s} \quad (4.3)$$

$$B^* = [1 + \sigma'_i; \rho_i^{(1,1)}, \dots, \rho_i^{(1,r)}, \tau_i^{(1,1)}, \dots, \tau_i^{(1,l)}, 0, \dots, 0]_{1,s}, \dots,$$

$$[1 + \sigma_i^{(T)}; \rho_i^{(T,1)}, \dots, \rho_i^{(T,r)}, \tau_i^{(T,1)}, \dots, \tau_i^{(T,l)}, 0, \dots, 0]_{1,s}, [1 - B_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0]_{1,Q},$$

$$[1 - \alpha_i - \beta_i; (\delta'_i + \eta'_i), \dots, (\delta_i^{(r)} + \eta_i^{(r)}), (\mu'_i + \theta'_i), \dots, (\mu_i^{(l)} + \theta_i^{(l)}), 1, \dots, 1]_{1,s} \quad (4.4)$$

The quantities $U, V, W, A, B; C$ and D are defined in the section I.

Formula 1

We have the following multiple Eulerian integral and we obtain the Aleph-function of $(r + l + T)$ -variables

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\mathfrak{N}_{U:W}^{0,n;V} \left(\begin{array}{c} z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta'_i} (v_i - x_i)^{\eta'_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,r)}}} \right] \end{array} \right)$$

$${}_P F_Q \left[(A_P); (B_Q); - \sum_{k=1}^l g_k \prod_{i=1}^s \left[\frac{(x_i - u_i)^{u_i^{(k)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,k)}}} \right] \right] dx_1 \cdots dx_s$$

$$= \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\mathfrak{N}_{U;sT+P+2s;V_1}^{0;n;sT+P+2s;W_1} \left(\begin{array}{c|c} z_1 w_1 & A ; A^* : C_1 \\ \cdots & \cdot \\ \cdots & \cdot \\ z_r w_r & \cdot \\ g_1 W_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ g_l W_l & \cdot \\ G_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ G_T & B ; B^* : D_1 \end{array} \right) \quad (4.5)$$

Where

$$w_m = \prod_{i=1}^s \left[(v_i - u_i)^{\delta_i^{(m)} + \eta_i^{(m)}} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{-\rho_i^{(j,m)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\rho_i^{(j,m)}} \right], m = 1, \dots, r \quad (4.6)$$

$$W_k = \prod_{i=1}^s \left[(v_i - u_i)^{\mu_i^{(k)} + \theta_i^{(k)}} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{-\tau_i^{(j,k)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{-\tau_i^{(j,k)}} \right], k = 1, \dots, l \quad (4.7)$$

$$G_j = \prod_{i=1}^s \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = 1, \dots, W \quad (4.8)$$

$$G_j = - \prod_{i=1}^s \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = W + 1, \dots, T \quad (4.9)$$

Provided that :

(A) $W \in [0, T]; u_i, v_i \in \mathbb{R}; i = 1, \dots, r$

(B) $\min\{\delta_i^{(t)}, \eta_i^{(t)}, \rho_i^{(j,t)}, \mu_i^{(k)}, \theta_i^{(k)}, \tau_i^{(j,k)}\} \geq 0; j = 1, \dots, T; i = 1, \dots, s; k = 1, \dots, l; t = 1, \dots, r$

(C) $\sigma_i^{(j)} \in \mathbb{R}, U_i^{(j)}, V_i^{(j)} \in \mathbb{C}, z_t, g_k \in \mathbb{C}; j = 1, \dots, s; t = 1, \dots, r; k = 1, \dots, l; t = 1, \dots, r$

(D) $\max \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = 1, \dots, W$ and

$$\max \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = W + 1, \dots, T$$

(E) $U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_i^{(k)}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)}$

$$- \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_i^{(k)}} \delta_{ji^{(k)}}^{(k)} \leq 0$$

(F) $A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_i^{(k)}} \gamma_{ji^{(k)}}^{(k)}$

$$+ \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_i^{(k)}} \delta_{ji^{(k)}}^{(k)} - \delta_i^{(k)} - \eta_i^{(k)} - \sum_{j=1}^T \rho_i^{(j,k)} > 0$$

with $k = 1, \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$

$$(G) \operatorname{Re} \left[\alpha_i + \sum_{t=1}^r \delta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{b_j^{(t)}}{\beta_j^{(t)}} \right] > 0; \operatorname{Re} \left[\beta_i + \sum_{t=1}^r \eta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{b_j^{(t)}}{\beta_j^{(t)}} \right] > 0; i = 1, \dots, s$$

$$(H) \left| \operatorname{arg} \left(z_i \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,k)}} \right) \right| < \frac{1}{2} A_i^{(k)} \pi$$

(I) See I

(J) $P \leq Q + 1$. The equality holds, when , in addition,

$$\text{either } P > Q \text{ and } \sum_{k=1}^l \left| g_k \left(\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,k)}} \right) \right|^{\frac{1}{Q-P}} < 1 \quad (u_i \leq x_i \leq v_i; i = 1, \dots, s)$$

$$\text{or } P \leq Q \text{ and } \max_{1 \leq k \leq l} \left[\left| \left(g_k \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\tau_i^{(j,k)}} \right) \right| \right] < 1 \quad (u_i \leq x_i \leq v_i; i = 1, \dots, s)$$

Proof

To establish the formula (3.5), we first use contour integral representation with the help of (1.2) for the multivariable Aleph-function occurring in its left-hand side and use the contour integral representation with the help of (3.1) for the generalized hypergeometric function ${}_P F_Q(\cdot)$. We write,

$$\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \quad (4.10)$$

$$\text{where } K_i^{(j)} = \eta_i^{(j)} - \sum_{t=1}^r \rho_i^{(j,t)} \zeta_t - \sum_{k=1}^l \tau_i^{(j,k)} \zeta_k \quad (4.11)$$

and express the factor occurring in R.H.S. Of (4.5) in terms of following Mellin-Barnes contour integral with the help of the result [9, page 18, eq.(2.6.4)]

$$\prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W \left[\frac{(U_i^{(j)} u_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^W} \int_{L'_1} \cdots \int_{L'_W} \prod_{j=1}^W \left[\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right] \prod_{j=1}^W \left[\frac{(U_i^{(j)} (x_i - u_i))^{\zeta'_j}}{(u_i U_i^{(j)} + V_i^{(j)})} \right] d\zeta'_1 \cdots d\zeta'_W \quad (4.12)$$

and

$$\prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=W+1}^T \left[\frac{(U_i^{(j)} v_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^{T-W}} \int_{L'_{W+1}} \cdots \int_{L'_{Tj=W+1}} \prod_{j=W+1}^T \left[\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right]$$

$$\prod_{j=W+1}^T \left[-\frac{(U_i^{(j)}(v_i - x_i))^{\zeta'_j}}{(v_i U_i^{(j)} + V_i^{(j)})} \right]^{\zeta'_j} d\zeta'_{W+1} \cdots d\zeta'_T \quad (4.13)$$

We apply the Fubini's theorem for multiple integral. Finally evaluating the innermost \mathbf{x} -integral with the help of (3.2) and reinterpreting the multiple Mellin-Barnes contour integral in terms of multivariable Aleph-function of $(r + l + T)$ -variables, we obtain the formula (4.5)

Formula 2

We note :

$$V_1 = V; 1, 1; 1, 0; \cdots; 1, 0; W_1 = W; 0, 1; \cdots; 0, 1 \quad (4.14)$$

$$C_1 = C; (1, 0); \cdots; (1, 0); D_1 = D; (0, 1), \cdots, (0, 1) \quad (4.15)$$

$$A^* = [1 + v'_i + \mu'_i K - \theta'_i R; \rho_i^{(1,1)}, \cdots, \rho_i^{(1,r)}, 1, 0, \cdots, 0]_{1,s}, \cdots,$$

$$[1 + v_i^{(T)} + \mu^{(T)} K - \theta_i^{(K)} R; \rho_i^{(T,1)}, \cdots, \rho_i^{(T,r)}, 1, 0, \cdots, 0]_{1,s},$$

$$[1 - \alpha_i - \epsilon_i K - \zeta_i R; \delta'_i, \cdots, \delta_i^{(r)}, 1, \cdots, 1, 0, \cdots, 0]_{1,s},$$

$$[1 - \beta_i - \sigma_i K - \lambda_i R; \eta'_i, \cdots, \eta_i^{(r)}, 0, \cdots, 0, 1, \cdots, 1]_{1,s} \quad (4.16)$$

$$B^* = [1 + v'_i + \mu'_i K - \theta'_i R; \rho_i^{(1,1)}, \cdots, \rho_i^{(1,r)}, 0, \cdots, 0]_{1,s}, \cdots,$$

$$[1 + v_i^{(T)} + \mu^{(T)} K - \theta_i^{(K)} R; \rho_i^{(T,1)}, \cdots, \rho_i^{(T,r)}, 0, \cdots, 0]_{1,s},$$

$$[1 - \alpha_i - \beta_i - (\epsilon_i + \sigma_i) K - (\zeta_i + \lambda_i) R; (\delta'_i + \eta'_i), \cdots, (\delta_i^{(r)} + \eta_i^{(r)}), 1, \cdots, 1]_{1,s} \quad (4.17)$$

In this section we will note $R_n^{\alpha, \beta}[x; E, F, g, h; p, q; \gamma; \delta; e^{-sx^r}]$ by $R_n^{\alpha, \beta}(x)$. We have the following multiple Eulerian integral

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\mathfrak{N}_{U:W}^{0,n;V} \left(\begin{matrix} z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta'_i} (v_i - x_i)^{\eta'_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{matrix} \right) R_n^{\alpha, \beta} \left[\prod_{j=1}^s \left[\frac{(x_i - u_i)^{\zeta_i} (v_i - x_i)^{\lambda_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\theta_i^{(j)}}} \right] \right]$$

$$S_N^M \left[\prod_{j=1}^s (x_i - u_i)^{\epsilon_i} (v_i - x_i)^{\sigma_i} \prod_{j=1}^T \left(U_i^{(j)} x_i + V_i^{(j)} \right)^{\mu_i^{(j)}} \right] dx_1 \cdots dx_r$$

$$= \prod_{j=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{v_i^{(j)}} \prod_{j=W+1}^T \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{v_i^{(j)}} \right] \sum_{K=0}^{[N/M]} \sum_{w, v, u, t, e, k_1, k_2}$$

$$\psi'(w, v, u, t, e, k_1, k_2) \mathfrak{N}_{U; sT+2s; sT+s: W_1}^{0; n; sT+2s: V_1} \left(\begin{array}{c|c} z_1 w_1 & \mathbf{A}; \mathbf{A}^* : C_1 \\ \cdots & \cdot \\ \cdots & \cdot \\ z_r w_r & \cdot \\ \cdots & \cdot \\ G_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ G_T & \mathbf{B}; \mathbf{B}^* : D_1 \end{array} \right) \quad (4.18)$$

where $\psi'(w, v, u, t, e, k_1, k_2)$

$$= \frac{(-n)_{MK} A_{N,K} \psi(w, v, u, t, e, k_1, k_2) \prod_{i=1}^s (v_i - u_i)^{(\epsilon_i + \sigma_i)K + (\zeta_i + \lambda_i)R}}{K! \prod_{i=1}^s \left[\prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{-\mu_i^{(j)}K + \theta_i^{(j)}R} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{-\mu_i^{(j)}K + \theta_i^{(j)}R} \right]} \quad (4.19)$$

$\psi(w, v, u, t, e, k_1, k_2)$ and R are given by (2.4) and (2.3) respectively.

$$w_l = \prod_{i=1}^s \left[(v_i - u_i)^{\delta_i^{(l)} + \eta_i^{(l)}} \prod_{j=1}^W \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-\rho_i^{(j,l)}} \prod_{j=W+1}^T \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{\rho_i^{(j,l)}} \right], l = 1, \dots, r \quad (4.20)$$

$$G_j = \prod_{i=1}^s \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = 1, \dots, W \quad (4.21)$$

$$G_j = - \prod_{i=1}^s \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = W + 1, \dots, T \quad (4.22)$$

Provided that :

(A) $W \in [0, T]; u_i, v_i \in \mathbb{R}; i = 1, \dots, r$

(B) $\min\{\zeta_i, \lambda_i, \theta_i^{(j)}, \delta_i^{(l)}, \eta_i^{(l)}, \rho_i^{(j,l)}\} \geq 0; j = 1, \dots, T; i = 1, \dots, s; l = 1, \dots, r$

(C) $\operatorname{Re}(\alpha_i, \beta_i, v_i^{(j)}) > 0 (i = 1, \dots, s; j = 1, \dots, T); |\tau| < 1$

$$(D) \max \left[\frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = 1, \dots, W \text{ and}$$

$$\max \left[\frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = W + 1, \dots, T$$

$$(E) U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i(k)} \sum_{j=n_k+1}^{p_i(k)} \gamma_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i(k)} \sum_{j=m_k+1}^{q_i(k)} \delta_{ji}^{(k)} \leq 0$$

$$(F) A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i(k)} \sum_{j=n_k+1}^{p_i(k)} \gamma_{ji}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i(k)} \sum_{j=m_k+1}^{q_i(k)} \delta_{ji}^{(k)} - \delta_i^{(k)} - \eta_i^{(k)} - \sum_{j=1}^T \rho_i^{(j,k)} > 0$$

$$(i = 1, \dots, s; k = 1, \dots, r)$$

$$(G) \operatorname{Re} \left[\alpha_i + \sum_{t=1}^r \delta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{b_j^{(t)}}{\beta_j^{(t)}} \right] > 0; \operatorname{Re} \left[\beta_i + \sum_{t=1}^r \eta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{b_j^{(t)}}{\beta_j^{(t)}} \right] > 0; i = 1, \dots, s$$

$$(H) \left| \arg \left(z_i \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,k)}} \right) \right| < \frac{1}{2} A_i^{(k)} \pi$$

(I) See I

(J) The series occurring on the right-hand side of (5.13) is absolutely and uniformly convergent

Proof

To establish the formula (4.18), we first use series representation (2.1) and (1.11) for $R_n^{\alpha, \beta}[\cdot]$ and $S_N^M(\cdot)$ respectively and contour integral representation with the help of (1.2) for the multivariable Aleph-function occurring in its left-hand side. Changing the order of integration and summation (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process). We have :

$$\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \quad (4.23)$$

$$\text{where } K_i^{(j)} = v_i^{(j)} - \theta_i^{(j)} R - \sum_{l=1}^r \rho_i^{(j,l)} \psi_l; i = 1, \dots, s; j = 1, \dots, T \quad (4.24)$$

and express the factors occurring in R.H.S. Of (4.18) in terms of following Mellin-Barnes contour integral, we obtain :

$$\prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W \left[\frac{(U_i^{(j)} u_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^W} \int_{L'_1} \cdots \int_{L'_W} \prod_{j=1}^W \left[\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right. \\ \left. \prod_{j=1}^W \left[\frac{(U_i^{(j)}(x_i - u_i))^{K_i^{(j)}}}{(u_i U_i^{(j)} + V_i^{(j)})} \right]^{\zeta'_j} \right] d\zeta'_1 \cdots d\zeta'_W \quad (4.25)$$

and

$$\prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=W+1}^T \left[\frac{(U_i^{(j)} v_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^{T-W}} \int_{L'_{W+1}} \cdots \int_{L'_T} \prod_{j=W+1}^T \left[\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right. \\ \left. \prod_{j=W+1}^T \left[\frac{(U_i^{(j)}(x_i - v_i))^{K_i^{(j)}}}{(v_i U_i^{(j)} + V_i^{(j)})} \right]^{\zeta'_j} \right] d\zeta'_{W+1} \cdots d\zeta'_T \quad (4.26)$$

We apply the Fubini's theorem for multiple integral. Finally evaluating the innermost \mathbf{x} -integral with the help of (3.2) and reinterpreting the multiple Mellin-Barnes contour integral in terms of multivariable I-function of $(r + T)$ -variables, we obtain the formula (4.18)

4. Multivariable I-function

If $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$, the multivariable Aleph-functions degenerate in multivariable I-function defined by Sharma and al [7] and we have two general multiple Eulerian integrals of multivariable I-functions.

Formula 3

We have the following multiple Eulerian integral and we obtain the I-function of $(r + l + T)$ -variables

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right] \\ I_{U:W}^{0, n:V} \left(\begin{matrix} Z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i^{(1)}} (v_i - x_i)^{\eta_i^{(1)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,1)}}} \right] \\ \vdots \\ Z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,r)}}} \right] \end{matrix} \right) \\ {}_P F_Q \left[(A_P); (B_Q); - \sum_{k=1}^l g_k \prod_{i=1}^s \left[\frac{(x_i - u_i)^{u_i^{(k)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,k)}}} \right] \right] dx_1 \cdots dx_s$$

$$\begin{aligned}
&= \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \right] \\
&I_{U; sT+P+2s; sT+Q+s; W_1}^{0; n; sT+P+2s; V_1} \left(\begin{array}{c|c} z_1 w_1 & A ; A^* : C_1 \\ \cdots & \cdot \\ \cdots & \cdot \\ z_r w_r & \cdot \\ g_1 W_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ g_l W_l & \cdot \\ G_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ G_T & B ; B^* : D_1 \end{array} \right) \tag{4.1}
\end{aligned}$$

under the same conditions and notations that (4.5) with $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$

Formula 4

We have the I-function of $(r + T)$ -variables

$$\begin{aligned}
&\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right] \\
&I_{U; W}^{0, n; V} \left(\begin{array}{c} z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i'} (v_i - x_i)^{\eta_i'}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \cdot \\ \cdot \\ z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{array} \right) R_n^{\alpha, \beta} \left[\prod_{j=1}^s \left[\frac{(x_i - u_i)^{\zeta_i} (v_i - x_i)^{\lambda_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\theta_i^{(j)}}} \right] \right]
\end{aligned}$$

$$S_N^M \left[\prod_{j=1}^s (x_i - u_i)^{\epsilon_i} (v_i - x_i)^{\sigma_i} \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\mu_i^{(j)}} \right] dx_1 \cdots dx_r$$

$$= \prod_{j=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{v_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{v_i^{(j)}} \right] \sum_{K=0}^{[N/M]} \sum_{w, v, u, t, e, k_1, k_2}$$

$$\psi'(w, v, u, t, e, k_1, k_2) I_{U; sT+2s; sT+s: W_1}^{0; n; sT+2s: V_1} \left(\begin{array}{c|c} z_1 w_1 & A ; A^* : C_1 \\ \cdots & \cdot \\ \cdots & \cdot \\ z_r w_r & \cdot \\ \cdots & \cdot \\ G_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ G_T & B ; B^* : D_1 \end{array} \right) \quad (4.2)$$

under the same conditions and notations that (4.18) with $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$

Remarks:

If $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$ and $R = R^{(1)} = \dots = R^{(r)} = 1$, the multivariable Aleph-function degenerates in multivariable H-function defined by Srivastava et al [11]. The formula (3.5) have been established by Goyal et al [4] and the formula (3.18) have been established by Garg [3].

5. Conclusion

In this paper we have evaluated two generalized multiple Eulerian integrals involving the multivariable Aleph-functions and multivariable I-function defined by Sharma et [7] with general arguments. The formulae established in this paper is very general nature. Thus, the results established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

[1] Agrawal B.D. And Chaubey J.P. Certain derivation of generating relations for generalized polynomials. Indian J. Pure and Appl. Math 10 (1980), page 1155-1157, ibid 11 (1981), page 357-359

[2] Fujiwara I. A unified presentation of classical orthogonal polynomials. Math. Japon. 11 (1966), page 133-148.

[3] Garg R. Unified multiple Eulerian integrals. Ganita Sandesh. Vol 15 (no 2) (2001), page 77-90.

[4] Goyal S.P. And Mathur T. On general multiple Eulerian integrals and fractional integration. Vijnana Parishad Anusandhan Patrika Vol 46 (no 3) , 2003 page 231-245.

[5] Raizada S.K. A study of unified representation of special functions of Mathematics Physics and their use in statistical and boundary value problem. Ph.D. Thesis, Bundelkhand University, Jhansi, India, 1991

[6] Salim T.O. A serie formula of generalized class of polynomials associated with Laplace transform and fractional integral operators. J. Rajasthan. Acad. Phy. Sci. 1(3) (2002), page 167-176.

[7] Sharma C.K. and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20, no2, p 113-116.

[8] Srivastava H.M., A contour integral involving Fox's H-function. Indian J.Math. 14(1972), page 1-6

[9] Srivastava H.M. Gupta K.C. And Goyal S.P. The H-functions of one and two variables with applications. South Asian Publishers Pvt Ltd 1982

[10] Srivastava H.M. and Karlsson P.W. Multiple Gaussian Hypergeometric series. Ellis.Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.

[11] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.

