

On general multiple Eulerian integrals involving the modified multivariable

H-function, a general class of polynomials and S generalized

Gauss hypergeometric function

F.Y. AYANT¹

¹ Teacher in High School , France
 E-mail : fredericayant@gmail.com

ABSTRACT

Goyal and Mathur [4], Garg [3] have studied the unified multiple Eulerian integrals. The object of this paper is first, to evaluate a general multiple Eulerian integrals with general integrands involving the product of a modified multivariable H-function defined by Prasad and Singh [5], a general Class of polynomials, the S generalized Gauss hypergeometric function and generalized hypergeometric function with general arguments. The second multiple Eulerian integral contain a general class of polynomials, the general polynomial set, the S generalized Gauss hypergeometric function and modified multivariable H-function defined by Prasad and Singh with general arguments. Our integral formulas are interesting and unified nature.

Keywords :Modified multivariable H-function, class of polynomial, general polynomials set, multiple Eulerian integral, S Generalized Gauss hypergeometric function, multivariable H-function, Srivastava-Daoust function.

2010 Mathematics Subject Classification. 33C60, 82C31

1. Introduction

In this paper, we evaluate two multiple Eulerian integrals involving the modified multivariable H-function defined by Prasad and Singh [5], the S generalized Gauss hypergeometric function and multivariable class of polynomials with general arguments. The modified H-function defined by Prasad and Singh [5] generalizes the multivariable H-function defined by Srivastava and Panda [11,13]. It is defined in term of multiple Mellin-Barnes type integral :

$$H(z_1, \dots, z_r) = H_{\mathbf{p}, \mathbf{q}; |R: p_1, q_1; \dots, p_r, q_r}^{\mathbf{m}, \mathbf{n}; |R': m_1, n_1; \dots, m_r, n_r} \left(\begin{matrix} Z_1 \\ \cdot \\ \cdot \\ \cdot \\ Z_r \end{matrix} \middle| \begin{matrix} (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, \mathbf{p}} : \\ \\ \\ (b_j; \beta_j^{(1)}, \dots, \beta_j^{(r)})_{1, \mathbf{q}} : \end{matrix} \right)$$

$$\left(\begin{matrix} (e_j; u_j' g_j', \dots, u_j^{(r)} g_j^{(r)})_{1, R'} : (c_j'; \gamma_j')_{1, p_1}, \dots, (c_j^{(r)}; \gamma_j^{(r)})_{1, p_r} \\ \\ (l_j; U_j' f_j', \dots, U_j^{(r)} f_j^{(r)})_{1, R} : (d_j'; \delta_j')_{1, q_1}, \dots, (d_j^{(r)}; \delta_j^{(r)})_{1, q_r} \end{matrix} \right) \quad (1.1)$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \phi(s_1, \dots, s_r) \prod_{i=1}^r \theta_i(s_i) z_i^{s_i} ds_1 \dots ds_r \quad (1.2)$$

where $\phi(s_1, \dots, s_r), \theta_i(s_i), i = 1, \dots, r$ are given by :

$$\phi(s_1, \dots, s_r) = \frac{\prod_{j=1}^m \Gamma(b_j - \sum_{i=1}^r \beta_j^{(i)} s_i) \prod_{j=1}^n \Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(i)} s_j)}{\prod_{j=n+1}^p \Gamma(a_j - \sum_{i=1}^r \alpha_j^{(i)} s_j) \prod_{j=m+1}^q \Gamma(1 - b_j + \sum_{i=1}^r \beta_j^{(i)} s_j)}$$

$$\frac{\prod_{j=1}^{R'} \Gamma(e_j + \sum_{i=1}^r u_j^{(i)} g_j^{(i)} s_i)}{\prod_{j=1}^R \Gamma(l_j + \sum_{i=1}^r U_j^{(i)} f_j^{(i)} s_i)} \quad (1.3)$$

$$\phi_i(s_i) = \frac{\prod_{j=1}^{n_i} \Gamma(1 - c_j^{(i)} + \gamma_j^{(i)} s_i) \prod_{j=1}^{m_i} \Gamma(d_j^{(i)} - \delta_j^{(i)} s_i)}{\prod_{j=n_i+1}^{p_i} \Gamma(c_j^{(i)} - \gamma_j^{(i)} s_i) \prod_{j=m_i+1}^{q_i} \Gamma(1 - d_j^{(i)} + \delta_j^{(i)} s_i)} \quad (1.4)$$

The multiple integral (1.17) converges absolutely if

$$|\arg Z_k| < \frac{1}{2} U_i \pi \quad (i = 1, \dots, r) \quad (1.5)$$

$$U_i = \sum_{j=1}^{\mathbf{m}} \beta_j^{(i)} - \sum_{j=\mathbf{m}+1}^{\mathbf{q}} \beta_j^{(i)} + \sum_{j=1}^{\mathbf{n}} \alpha_j^{(i)} - \sum_{j=\mathbf{n}+1}^{\mathbf{p}} \alpha_j^{(i)} + \sum_{j=1}^{m_i} \delta_j^{(i)} - \sum_{j=1+m_i}^{q_i} \delta_j^{(i)} \\ + \sum_{j=1}^{n_i} \gamma_j^{(i)} - \sum_{j=n_i+1}^{p_i} \gamma_j^{(i)} + \sum_{j=1}^{R'} g_j^{(i)} - \sum_{j=1}^R f_j^{(i)} > 0 \quad (i = 1, \dots, r) \quad (1.6)$$

Srivastava and Garg [10] introduced and defined a general class of multivariable polynomials as follows

$$S_L^{h_1, \dots, h_u} [z_1, \dots, z_u] = \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{z_1^{R_1} \dots z_u^{R_u}}{R_1! \dots R_u!} \quad (1.7)$$

The coefficients $B(E; R_1, \dots, R_u)$ are arbitrary constants, real or complex.

We shall note

$$B_u = \frac{(-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u)}{R_1! \dots R_u!} \quad (1.8)$$

2. Sequence of functions

Agarwal and Chaubey [1], Salim [7] and several others have studied a general sequence of functions. In the present document we shall study the following useful series formula for a general sequence of functions.

$$R_n^{\alpha, \beta} [x; E, F, g, h; p, q; \gamma; \delta; e^{-s x^r}] = \sum_{w, v, u, t, e, k_1, k_2} \psi(w, v, u, t, e, k_1, k_2) x^R \quad (2.1)$$

$$\text{where } \sum_{w, v, u, t, e, k_1, k_2} = \sum_{w=0}^{\infty} \sum_{v=0}^n \sum_{u=0}^v \sum_{t=0}^n \sum_{c=0}^t \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \quad (2.2)$$

and the infinite series on the right side (2.1) is absolutely convergent, $R = ln + qv + pt + rw + k_1 r + k_2 q$ (2.3)

$$\text{and } \psi(w, v, u, t, e, k_1, k_2) = \frac{(-)^{t+w+k_2} (-v)_u (-t)_e (\alpha)_t l^n s^{w+k_1} F^{\gamma n-t}}{w! v! u! t! e! K_n k_1! k_2!} \frac{s^{w+k_1} F^{\gamma n-t}}{(1 - \alpha - t)_e} (\alpha - \gamma n)_e$$

$$(-\beta - \delta n)_v g^{v+k_2} h^{\delta n-v-k_2} (v - \delta n)_{k_2} E^t \left(\frac{pe + rw + \lambda + qn}{l} \right)_n \quad (2.4)$$

where K_n is a sequence of constants. This function will note $R_n^{\alpha, \beta}[x]$

By suitably specializing the parameters involving in (2.1), a general sequence of function reduced to generalized polynomials set studied by Raizada [6], a class of polynomials introduced by Fujiwara [2] and several others authors.

3. S Generalized Gauss's hypergeometric function

The S generalized Gauss hypergeometric function $F_p^{(\alpha, \beta; \tau, \mu)}(a, b, c; z)$ introduced and defined by Srivastava et al [8, page 350, Eq.(1.12)] is represented in the following manner :

$$F_p^{(\alpha, \beta; \tau, \mu)}(a, b, c; z) = \sum_{n=0}^{\infty} (a)_n \frac{B_p^{(\alpha, \beta; \tau, \mu)}(b+n, c-b)}{B(b, c-b)} \frac{z^n}{n!} \quad (|z| < 1) \quad (3.1)$$

provided that $(Re(p) \geq 0, \min Re(\alpha, \beta, \tau, \mu) > 0; Re(c) > Re(b) > 0)$

where the S generalized Beta function $B_p^{(\alpha, \beta; \tau, \mu)}(x, y)$ was introduced and defined by Srivastava et al [8, page 350, Eq(1.13)]

$$B_p^{(\alpha, \beta; \tau, \mu)}(x, y) = \int_1^1 t^{x-1} (1-t)^{y-1} {}_1F_1 \left(\alpha; \beta; \frac{-p}{t^\tau (1-t)^\mu} \right) dt \quad (3.2)$$

provided that $(Re(p) \geq 0, \min Re(x, y, \alpha, \beta) > 0; \min\{Re(\tau), Re(\mu)\} > 0)$

4. Integral representation of generalized hypergeometric function

The following generalized hypergeometric function in terms of multiple contour integrals is also required [12, page 39 eq .30]

$$\begin{aligned} & \frac{\prod_{j=1}^P \Gamma(A_j)}{\prod_{j=1}^Q \Gamma(B_j)} {}_P F_Q [(A_P); (B_Q); -(x_1 + \dots + x_r)] \\ &= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \frac{\prod_{j=1}^P \Gamma(A_j + s_1 + \dots + s_r)}{\prod_{j=1}^Q \Gamma(B_j + s_1 + \dots + s_r)} \Gamma(-s_1) \dots \Gamma(-s_r) x_1^{s_1} \dots x_r^{s_r} ds_1 \dots ds_r \end{aligned} \quad (4.1)$$

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of $\Gamma(A_j + s_1 + \dots + s_r)$ are separated from those of $\Gamma(-s_j), j = 1, \dots, r$. The above result (1.23) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of $\Gamma(-s_j), j = 1, \dots, r$

The equivalent form of Eulerian beta integral is :

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} dt = (b-a)^{\alpha+\beta} B(\alpha, \beta) (Re(\alpha) > 0, Re(\beta) > 0, a \neq b) \quad (4.2)$$

5. First integral

We shall note :

$$X = m_1, n_1; \cdots; m_r, n_r; 1, 0; \cdots; 1, 0; 1, 0; \cdots; 1, 0 \quad (5.1)$$

$$Y = p_1, q_1; \cdots; p_r, q_r; 0, 1; \cdots; 0, 1; 0, 1; \cdots; 0, 1 \quad (5.2)$$

$$\mathbb{A} = (a_j; \alpha_j^{(1)}, \cdots, \alpha_j^{(r)}, 0, \cdots, 0, 0, \cdots, 0)_{1,p} : (e_j; u'_j g'_j, \cdots, u_j^{(r)} g_j^{(r)}; 0, \cdots, 0, 0, \cdots, 0)_{1,R'} : \\ (c'_j; \gamma'_j)_{1,p_1}, \cdots, (c_j^{(r)}; \gamma_j^{(r)})_{1,p_r}; (1, 0); \cdots; (1, 0); (1, 0); \cdots; (1, 0) \quad (5.3)$$

$$\mathbb{B} = (b_j; \beta_j^{(1)}, \cdots, \beta_j^{(r)}, 0, \cdots, 0, 0, \cdots, 0)_{1,q} : (l_j; U'_j f'_j, \cdots, U_j^{(r)} f_j^{(r)}, 0, \cdots, 0, 0, \cdots, 0)_{1,R} : \\ (d'_j; \delta'_j)_{1,q_1}, \cdots, (d_j^{(r)}; \delta_j^{(r)})_{1,q_r}; (0, 1); \cdots; (0, 1); (0, 1); \cdots; (0, 1) \quad (5.4)$$

$$A^* = [1 + \sigma'_i - n'c'_i - \sum_{k=1}^u R_k \xi_i^{(1,k)}; \rho_i^{(1,1)}, \cdots, \rho_i^{(1,r)}, \tau_i^{(1,1)}, \cdots, \tau_i^{(1,l)}, 1, 0, \cdots, 0]_{1,s}, \cdots,$$

$$[1 + \sigma_i^{(T)} - n'c_i^{(T)} - \sum_{k=1}^u R_k \xi_i^{(T,k)}; \rho_i^{(T,1)}, \cdots, \rho_i^{(T,r)}, \tau_i^{(T,1)}, \cdots, \tau_i^{(T,l)}, 0, \cdots, 0, 1]_{1,s},$$

$$[1 - A_j; 0, \cdots, 0, 1, \cdots, 1, 0, \cdots, 0]_{1,P},$$

$$[1 - \alpha_i - n'a_i - \sum_{k=1}^u R_k \alpha_i^{(k)}; \delta_i^{(1)}, \cdots, \delta_i^{(r)}, \mu'_i, \cdots, \mu_i^{(l)}, 1, \cdots, 1, 0, \cdots, 0]_{1,s},$$

W-items (T-W)-items

$$[1 - \beta_i - n'b_i - \sum_{k=1}^u R_k \beta_i^{(k)}; \eta'_i, \cdots, \eta_i^{(r)}, \theta'_i, \cdots, \theta_i^{(l)}, 0, \cdots, 0, 1, \cdots, 1]_{1,s} \quad (5.5)$$

W-items (T-W)-items

$$B^* = [1 + \sigma'_i - n'c'_i - \sum_{k=1}^u R_k \xi_i^{(1,k)}; \rho_i^{(1,1)}, \cdots, \rho_i^{(1,r)}, \tau_i^{(1,1)}, \cdots, \tau_i^{(1,l)}, 0, \cdots, 0]_{1,s}, \cdots,$$

$$[1 + \sigma_i^{(T)} - n'c_i^{(T)} - \sum_{k=1}^u R_k \xi_i^{(T,k)}; \rho_i^{(T,1)}, \cdots, \rho_i^{(T,r)}, \tau_i^{(T,1)}, \cdots, \tau_i^{(T,l)}, 0, \cdots, 0]_{1,s},$$

$$[1 - B_j; 0, \cdots, 0, 1, \cdots, 1, 0, \cdots, 0]_{1,Q},$$

$$[1 - \alpha_i - \beta_i - n'(a_i + b_i) - \sum_{k=1}^u R_k (\alpha_i^{(k)} + \beta_i^{(k)}); (\delta_i^{(1)} + \eta'_i), \cdots, (\delta_i^{(r)} + \eta_i^{(r)})$$

$$(\mu'_i + \theta'_i), \cdots, (\mu_i^{(l)} + \theta_i^{(l)}), 1, \cdots, 1]_{1,s} \quad (5.6)$$

We have the following multiple Eulerian integral and we obtain the modified H-function of $(r + l + T)$ -variables

Theorem 1

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$F_p^{(\alpha, \beta; \tau, \mu)} \left[a, b, c; z \prod_{i=1}^s \left[\frac{(x_i - u_i)^{a_i} (v_i - x_i)^{b_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{c_i^{(j)}}} \right] \right]$$

$$S_L^{h_1, \dots, h_u} \left(\begin{array}{c} y_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(1)}} (v_i - x_i)^{\beta_i'^{(1)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,1)}}} \right] \\ \vdots \\ y_u \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(u)}} (v_i - x_i)^{\beta_i'^{(u)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,u)}}} \right] \end{array} \right)$$

$$H \left(\begin{array}{c} z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i'^{(1)}} (v_i - x_i)^{\eta_i'^{(1)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i'^{(r)}} (v_i - x_i)^{\eta_i'^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{array} \right)$$

$${}_p F_Q \left[(A_P); (B_Q); - \sum_{k=1}^l g_k \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\mu_i^{(k)}} (v_i - x_i)^{\theta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,k)}}} \right] \right] dx_1 \cdots dx_s$$

$$= \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{i=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\sum_{n'=0}^{\infty} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} B_u(a)_{n'} \frac{B_p^{(\alpha, \beta; \tau, \mu)}(b + n', c - b)}{B(b, c - b)} \frac{z^{n'}}{n'!} y_1^{R_1} \cdots y_u^{R_u} \prod_{i=1}^s \left[(v_i - u_i)^{n'(a_i + b_i) + \sum_{k=1}^u R_k (\alpha_i'^{(k)} + \beta_i'^{(k)})} \right]$$

$$\prod_{i=1}^s \left[\prod_{j=1}^W \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)}} \prod_{j=W+1}^T \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)}} \right]$$

$$H_{\mathbf{p}+sT+P+2s, \mathbf{q}+sT+Q+s; |R:Y}^{\mathbf{m}, \mathbf{n}+sT+P+2s; |R':X} \left(\begin{array}{c|c} z_1 w_1 & \mathbb{A}, \mathbb{A}^* \\ \cdots & \cdot \\ \cdots & \cdot \\ z_r w_r & \cdot \\ g_1 W_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ g_l W_l & \cdot \\ G_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ G_T & \mathbb{B}, \mathbb{B}^* \end{array} \right) \quad (5.7)$$

Where

$$w_m = \prod_{i=1}^s \left[(v_i - u_i)^{\delta_i^{(m)} + \eta_i^{(m)}} \prod_{j=1}^W \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-\rho_i^{(j,m)}} \prod_{j=W+1}^T \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-\rho_i^{(j,m)}} \right], m = 1, \dots, r \quad (5.8)$$

$$W_k = \prod_{i=1}^s \left[(v_i - u_i)^{\mu_i^{(k)} + \theta_i^{(k)}} \prod_{j=1}^W \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-\tau_i^{(j,k)}} \prod_{j=W+1}^T \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-\tau_i^{(j,k)}} \right], k = 1, \dots, l \quad (5.9)$$

$$G_j = \prod_{i=1}^s \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = 1, \dots, W \quad (5.10)$$

$$G_j = - \prod_{i=1}^s \left[\frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = W + 1, \dots, T \quad (5.11)$$

Provided that :

(A) $0 \leq W \leq T; u_i, v_i \in \mathbb{R}; \min\{a_i, b_i, c_i^{(j)}, \alpha_i^{(k)}, \beta_i^{(k)}, \xi_i^{(j,k)}\} > 0, i = 1, \dots, s; k = 1, \dots, u; j = 1, \dots, T$

(B) $\min\{\delta_i^{(t)}, \eta_i^{(t)}, \rho_i^{(j,t)}, \mu_i^{(k)}, \theta_i^{(k)}, \tau_i^{(j,k)}\} \geq 0; j = 1, \dots, T; i = 1, \dots, s; k = 1, \dots, l; t = 1, \dots, r$

(C) $\sigma_i^{(j)} \in \mathbb{R}, U_i^{(j)}, V_i^{(j)} \in \mathbb{C}, z_t, g_k \in \mathbb{C}; j = 1, \dots, s; t = 1, \dots, r; k = 1, \dots, l; t = 1, \dots, r$

(D) $\max \left[\left| \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right| \right] < 1, i = 1, \dots, s; j = 1, \dots, W$ and

$$\max \left[\left| \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right| \right] < 1, i = 1, \dots, s; j = W + 1, \dots, T$$

$$\begin{aligned}
\text{(E)} \quad U'_i &= \sum_{j=1}^{\mathbf{m}} \beta_j^{(i)} - \sum_{j=\mathbf{m}+1}^{\mathbf{q}} \beta_j^{(i)} + \sum_{j=1}^{\mathbf{n}} \alpha_j^{(i)} - \sum_{j=\mathbf{n}+1}^{\mathbf{p}} \alpha_j^{(i)} + \sum_{j=1}^{m_i} \delta_j^{(i)} - \sum_{j=1+m_i}^{q_i} \delta_j^{(i)} \\
&+ \sum_{j=1}^{n_i} \gamma_j^{(i)} - \sum_{j=n_i+1}^{p_i} \gamma_j^{(i)} + \sum_{j=1}^{R'} g_j^{(i)} - \sum_{j=1}^R f_j^{(i)} - \delta_i^{(k)} - \eta_i^{(k)} - \sum_{j=1}^T \rho_i^{(j,k)} > 0
\end{aligned}$$

$$(i = 1, \dots, s; k = 1, \dots, r)$$

$$\text{(F)} \quad \operatorname{Re}(\alpha_i + n' a_i) + \sum_{t=1}^r \delta_i^{(t)} \min_{1 \leq j \leq m_i} \operatorname{Re} \left(\frac{d_j^{(t)}}{\delta_j^{(t)}} \right) > 0 \text{ and}$$

$$\operatorname{Re}(\beta_i + n' b_i) + \sum_{t=1}^r \eta_i^{(t)} \min_{1 \leq j \leq m_i} \operatorname{Re} \left(\frac{d_j^{(t)}}{\delta_j^{(t)}} \right) > 0; i = 1, \dots, s$$

$$\text{(G)} \quad \left| \arg \left(z_i \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\rho_i^{(j,k)}} \right) \right| < \frac{1}{2} U'_i \pi$$

(H) $P \leq Q + 1$. The equality holds, when , in addition,

$$\text{either } P > Q \text{ and } \sum_{k=1}^l \left| g_k \left(\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\tau_i^{(j,k)}} \right) \right|^{\frac{1}{Q-P}} < 1 \quad (u_i \leq x_i \leq v_i; i = 1, \dots, s)$$

$$\text{or } P \leq Q \text{ and } \max_{1 \leq k \leq l} \left[\left| \left(g_k \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\tau_i^{(j,k)}} \right) \right| \right] < 1 \quad (u_i \leq x_i \leq v_i; i = 1, \dots, s)$$

(I) $(\operatorname{Re}(p) \geq 0, \min \operatorname{Re}(x, y, \alpha, \beta) > 0; \operatorname{Min}\{\operatorname{Re}(\tau), \operatorname{Re}(\mu)\} > 0)$

(J) The series occuring on the right-hand side of (5.13) are absolutely and uniformly convergent

Proof

To establish the formula (5.7), we first use series representation (3.1) and (1.7) for $F_p^{(\alpha, \beta; \tau, \mu)}[\cdot]$ and $S_L^{h_1, \dots, h_u}[\cdot]$ respectively, we use contour integral representation with the help of (1.1) for the modified multivariable H-function occuring in its left-hand side and use the contour integral representation with the help of (4.1) for the generalized hypergeometric function ${}_P F_Q(\cdot)$. Changing the order of integration and summation (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process). Now we write :

$$\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}}$$

$$\text{where } K_i^{(j)} = \eta_i^{(j)} - n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)} - \sum_{t=1}^r \rho_i^{(j,t)} \zeta_t - \sum_{k=1}^l \tau_i^{(j,k)} \zeta_k \quad i = 1, \dots, s; j = 1, \dots, T \quad (5.12)$$

and express the factor occuring in R.H.S. Of (5.13) in terms of following Mellin-Barnes contour integral with the help of the result [11, page 18, eq.(2.6.4)].

$$\prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W \left[\frac{(U_i^{(j)} u_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^W} \int_{L'_1} \cdots \int_{L'_W} \prod_{j=1}^W [\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j)] \prod_{j=1}^W \left[\frac{(U_i^{(j)} (x_i - u_i))^{K_i^{(j)}}}{(u_i U_i^{(j)} + V_i^{(j)})^{K_i^{(j)}}} \right] d\zeta'_1 \cdots d\zeta'_W \quad (5.13)$$

and

$$\prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=W+1}^T \left[\frac{(U_i^{(j)} v_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^{T-W}} \int_{L'_{W+1}} \cdots \int_{L'_{Tj=W+1}} \prod_{j=W+1}^T [\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j)] \prod_{j=W+1}^T \left[-\frac{(U_i^{(j)} (v_i - x_i))^{K_i^{(j)}}}{(v_i U_i^{(j)} + V_i^{(j)})^{K_i^{(j)}}} \right] d\zeta'_{W+1} \cdots d\zeta'_T \quad (5.14)$$

We apply the Fubini's theorem for multiple integral. Finally evaluating the innermost \mathbf{x} -integral with the help of (4.2) and reinterpreting the multiple Mellin-Barnes contour integral in terms of modified multivariable H-function of $(r + l + T)$ -variables, we obtain the formula (5.7).

6. Second formula

We shall note :

$$X = m_1, n_1; \cdots; m_r, n_r; 1, 0; \cdots; 1, 0 \quad (6.1)$$

$$Y = p_1, q_1; \cdots; p_r, q_r; 0, 1; \cdots; 0, 1 \quad (6.2)$$

$$\mathbb{A} = (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)}, 0, \dots, 0)_{1,p} : (e_j; u'_j g'_j, \dots, u_j^{(r)} g_j^{(r)}; 0, \dots, 0)_{1,R'} :$$

$$(c'_j; \gamma'_j)_{1,p_1}, \dots, (c'_j; \gamma'_j)_{1,p_r}; (1, 0); \cdots; (1, 0) \quad (6.3)$$

$$\mathbb{B} = (b_j; \beta_j^{(1)}, \dots, \beta_j^{(r)}, 0, \dots, 0)_{1,q} : (l_j; U'_j f'_j, \dots, U_j^{(r)} f_j^{(r)}, 0, \dots, 0)_{1,R} :$$

$$(d'_j; \delta'_j)_{1,q_1}, \dots, (d'_j; \delta'_j)_{1,q_r}; (0, 1); \cdots; (0, 1) \quad (6.4)$$

$$A^* = [1 + \sigma'_i - \theta'_i R - n' c'_i - \sum_{k=1}^u R_k \xi_i^{(1,k)}; \rho_i^{(1,1)}, \dots, \rho_i^{(1,r)}, 1, 0, \dots, 0]_{1,s}, \dots,$$

$$[1 + \sigma_i^{(T)} - \theta_i^{(T)} R - n' c_i^{(T)} - \sum_{k=1}^u R_k \xi_i^{(T,k)}; \rho_i^{(T,1)}, \dots, \rho_i^{(T,r)}, 0, \dots, 0, 1]_{1,s},$$

$$[1 - \alpha_i - \zeta_i R - n' a_i - \sum_{k=1}^u R_k \alpha_i^{(k)}; \delta_i^{(1)}, \dots, \delta_i^{(r)}, 1, \dots, 1, 0, \dots, 0]_{1,s},$$

W-items (T-W)-items

$$[1 - \beta_i - \lambda_i R - n' b_i - \sum_{k=1}^u R_k \beta_i^{(k)}; \eta_i', \dots, \eta_i^{(r)}, 0, \dots, 0, 1, \dots, 1]_{1,s} \quad (6.5)$$

W-items (T-W)-items

$$B^* = [1 + \sigma_i' - \theta_i' R - n' c_i' - \sum_{k=1}^u R_k \xi_i^{(1,k)}; \rho_i^{(1,1)}, \dots, \rho_i^{(1,r)}, 0, \dots, 0]_{1,s}, \dots,$$

$$[1 + \sigma_i^{(T)} - \theta_i^{(T)} R - n' c_i^{(T)} - \sum_{k=1}^u R_k \xi_i^{(T,k)}; \rho_i^{(T,1)}, \dots, \rho_i^{(T,r)}, 0, \dots, 0]_{1,s},$$

$$[1 - \alpha_i - \beta_i (\zeta_i + \lambda_i) R - n' (a_i + b_i) - \sum_{k=1}^u R_k (\alpha_i'^{(k)} + \beta_i'^{(k)}); (\delta_i^{(1)} + \eta_i'), \dots, (\delta_i^{(r)} + \eta_i^{(r)})$$

$$(\mu_i' + \theta_i'), \dots, (\mu_i^{(l)} + \theta_i^{(l)}), 1, \dots, 1]_{1,s} \quad (6.6)$$

We have the following multiple Eulerian integral

Theorem 2

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$F_p^{(\alpha, \beta; \tau, \mu)} \left[a, b, c; z \prod_{i=1}^s \left[\frac{(x_i - u_i)^{a_i} (v_i - x_i)^{b_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{c_i^{(j)}}} \right] \right] R_n^{\alpha, \beta} \left[Z \prod_{j=1}^s \left[\frac{(x_i - u_i)^{\zeta_i} (v_i - x_i)^{\lambda_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\theta_i^{(j)}}} \right] \right]$$

$$S_L^{h_1, \dots, h_u} \left(\begin{array}{c} y_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(1)}} (v_i - x_i)^{\beta_i'^{(1)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,1)}}} \right] \\ \vdots \\ y_u \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(u)}} (v_i - x_i)^{\beta_i'^{(u)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,u)}}} \right] \end{array} \right)$$

$$\begin{aligned}
& H \left(\begin{array}{c} z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta'_i(1)} (v_i - x_i)^{\eta'_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta'_i(r)} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{array} \right) dx_1 \cdots dx_s \\
&= \prod_{i=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\rho_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\rho_i^{(j)}} \right] \sum_{n'=0}^{\infty} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \sum_{w, v, u, t, e, k_1, k_2} \\
& B_u(a)_{n'} \frac{B_p^{(\alpha, \beta; \tau, \mu)}(b + n', c - b) z^{n'}}{B(b, c - b) n'!} y_1^{R_1} \cdots y_u^{R_u} \psi'(w, v, u, t, e, k_1, k_2) Z^R \prod_{i=1}^s \left[(v_i - u_i)^{n'(a_i + b_i) + \sum_{k=1}^u R_k (\alpha_i^{(k)} + \beta_i^{(k)})} \right] \\
& \prod_{i=1}^s \left[\prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{-n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{-n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)}} \right] \\
& \prod_{j=1}^s \left[(v_i - u_i)^{n'(a_i + b_i) + \sum_{k=1}^u R_k (\alpha_i^{(k)} + \beta_i^{(k)})} \right] H_{\mathbf{p} + sT + 2s, \mathbf{q} + sT + s; |R:Y}^{\mathbf{m}, \mathbf{n} + sT + 2s; |R:X} \left(\begin{array}{c|c} z_1 w_1 & \mathbb{A}, A^* \\ \vdots & \vdots \\ \vdots & \vdots \\ z_r w_r & \vdots \\ \vdots & \vdots \\ G_1 & \vdots \\ \vdots & \vdots \\ \vdots & \vdots \\ G_T & \mathbb{B}, B^* \end{array} \right) \quad (6.7)
\end{aligned}$$

where

$$\psi'(w, v, u, t, e, k_1, k_2) = \frac{\psi(w, v, u, t, e, k_1, k_2) \prod_{i=1}^s (v_i - u_i)^{(\zeta_i + \lambda_i)R}}{\prod_{i=1}^s \left[\prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\theta_i^{(j)} R} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\theta_i^{(j)} R} \right]} \quad (6.8)$$

$\psi(w, v, u, t, e, k_1, k_2)$ and R are given by (2.4) and (2.3) respectively.

$$w_l = \prod_{i=1}^s \left[(v_i - u_i)^{\delta_i^{(l)} + \eta_i^{(l)}} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{-\rho_i^{(j,l)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{-\rho_i^{(j,l)}} \right], l = 1, \dots, r \quad (6.9)$$

$$G_j = \prod_{i=1}^s \left[\frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = 1, \dots, W \quad (6.10)$$

$$G_j = - \prod_{i=1}^s \left[\frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = W + 1, \dots, T \quad (6.11)$$

Provided that :

$$(A) 0 \leq W \leq T; u_i, v_i \in \mathbb{R}; \min\{a_i, b_i, c_i^{(j)}, \alpha_i'^{(k)}, \beta_i'^{(k)}, \xi_i^{(j,k)}\} > 0, i = 1, \dots, s; k = 1, \dots, u; j = 1, \dots, T$$

$$(B) \min\{\zeta_i, \lambda_i, \theta_i^{(j)}, \delta_i^{(l)}, \eta_i^{(l)}, \rho_i^{(j,l)}\} \geq 0; j = 1, \dots, T; i = 1, \dots, s; l = 1, \dots, r$$

$$(C) \operatorname{Re}(\alpha_i, \beta_i, v_i^{(j)}) > 0 (i = 1, \dots, s; j = 1, \dots, T); |\tau| < 1$$

$$(D) \max \left[\left| \frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right| \right] < 1, i = 1, \dots, s; j = 1, \dots, W \text{ and}$$

$$\max \left[\left| \frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right| \right] < 1, i = 1, \dots, s; j = W + 1, \dots, T$$

$$(E) U_i' = \sum_{j=1}^m \beta_j^{(i)} - \sum_{j=m+1}^q \beta_j^{(i)} + \sum_{j=1}^n \alpha_j^{(i)} - \sum_{j=n+1}^p \alpha_j^{(i)} + \sum_{j=1}^{m_i} \delta_j^{(i)} - \sum_{j=1+m_i}^{q_i} \delta_j^{(i)}$$

$$+ \sum_{j=1}^{n_i} \gamma_j^{(i)} - \sum_{j=n_i+1}^{p_i} \gamma_j^{(i)} + \sum_{j=1}^{R'} g_j^{(i)} - \sum_{j=1}^R f_j^{(i)} - \delta_i'^{(k)} - \eta_i^{(k)} - \sum_{j=1}^T \rho_i^{(j,k)} > 0$$

$$(i = 1, \dots, s; k = 1, \dots, r)$$

$$(F) \operatorname{Re}(\alpha_i + n'a_i + R\zeta_i) + \sum_{t=1}^r \delta_i'^{(t)} \min_{1 \leq j \leq m_i} \operatorname{Re} \left(\frac{d_j^{(t)}}{\delta_j^{(t)}} \right) > 0 \text{ and}$$

$$\operatorname{Re}(\beta_i + n'b_i + R\lambda_i) + \sum_{t=1}^r \eta_i^{(t)} \min_{1 \leq j \leq m_i} \operatorname{Re} \left(\frac{d_j^{(t)}}{\delta_j^{(t)}} \right) > 0; i = 1, \dots, s$$

$$(G) \left| \arg \left(z_i \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\rho_i^{(j,k)}} \right) \right| < \frac{1}{2} U_i' \pi$$

(H) The series occurring on the right-hand side of (5.13) are absolutely and uniformly convergent

$$(I) (\operatorname{Re}(p) \geq 0, \min \operatorname{Re}(x, y, \alpha, \beta) > 0; \operatorname{Min}\{\operatorname{Re}(\tau), \operatorname{Re}(\mu)\} > 0)$$

Proof

To establish the formula (6.7), we first use series representation (3.1), (1.7) and (2.1) for $F_p^{(\alpha,\beta;\tau,\mu)}[\cdot]$, $S_L^{h_1,\dots,h_u}[\cdot]$ and $R_n^{\alpha,\beta}[\cdot]$ respectively and the contour integral representation with the help of (1.2) for the modified multivariable H-function defined by Prasad and Singh [5] occurring in its left-hand side. Changing the order of integration and summation (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process). Now, we write:

$$\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \quad (6.18)$$

$$\text{where } K_i^{(j)} = \eta_i^{(j)} - n' c_i^{(j)} - R \theta_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)} - \sum_{t=1}^r \rho_i^{(j,t)} \zeta_t; i = 1, \dots, s; j = 1, \dots, T \quad (6.19)$$

and express the factors occurring in R.H.S. Of (6.13) in terms of following Mellin-Barnes contour integral, we obtain :

$$\prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W \left[\frac{(U_i^{(j)} u_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^W} \int_{L'_1} \cdots \int_{L'_W} \prod_{j=1}^W \left[\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right. \\ \left. \prod_{j=1}^W \left[\frac{(U_i^{(j)} (x_i - u_i))^{\zeta'_j}}{(u_i U_i^{(j)} + V_i^{(j)})} \right] \right] d\zeta'_1 \cdots d\zeta'_W \quad (6.20)$$

and

$$\prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=W+1}^T \left[\frac{(U_i^{(j)} v_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^{T-W}} \int_{L'_{W+1}} \cdots \int_{L'_T} \prod_{j=W+1}^T \left[\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right. \\ \left. \prod_{j=W+1}^T \left[\frac{(U_i^{(j)} (x_i - v_i))^{\zeta'_j}}{(v_i U_i^{(j)} + V_i^{(j)})} \right] \right] d\zeta'_{W+1} \cdots d\zeta'_T \quad (6.21)$$

We apply the Fubini's theorem for multiple integral. Finally evaluating the innermost \mathbf{x} -integral with the help of (3.2) and reinterpreting the multiple Mellin-Barnes contour integral in terms of modified multivariable H-function of $(r + T)$ -variables, we obtain the formula (6.7)

7. Srivastava-daoust function

$$\text{If } B(L; R_1, \dots, R_u) = \frac{\prod_{j=1}^{\bar{A}} (a_j)_{R_1 \theta'_j + \dots + R_u \theta_j^{(u)}} \prod_{j=1}^{B'} (b'_j)_{R_1 \phi'_j} \cdots \prod_{j=1}^{B^{(u)}} (b_j^{(u)})_{R_u \phi_j^{(u)}}}{\prod_{j=1}^{\bar{C}} (c_j)_{m_1 \psi'_j + \dots + m_u \psi_j^{(u)}} \prod_{j=1}^{D'} (d'_j)_{R_1 \delta'_j} \cdots \prod_{j=1}^{D^{(u)}} (d_j^{(u)})_{R_u \delta_j^{(u)}}} \quad (7.1)$$

then the general class of multivariable polynomial $S_L^{h_1,\dots,h_u} [z_1, \dots, z_u]$ reduces to generalized Lauricella function defined by Srivastava et al [9].

$$F_{\bar{C}; D'; \dots; D^{(u)}}^{1+\bar{A}; B'; \dots; B^{(u)}} \left(\begin{array}{c} z_1 \\ \cdot \\ \cdot \\ z_u \end{array} \middle| \begin{array}{l} (-L; R_1, \dots, R_u), [(a); \theta', \dots, \theta^{(u)}] : [(b'); \phi']; \dots; [(b^{(u)}); \phi^{(u)}] \\ \\ [(c); \psi', \dots, \psi^{(u)}] : [(d'); \delta']; \dots; [(d^{(u)}); \delta^{(u)}] \end{array} \right) \quad (7.2)$$

and we have the two following formulae

Corollary 1

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$F_p^{(\alpha, \beta; \tau, \mu)} \left[a, b, c; z \prod_{i=1}^s \left[\frac{(x_i - u_i)^{a_i} (v_i - x_i)^{b_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{c_i^{(j)}}} \right] \right]$$

$$F_{\bar{C}:D'; \dots; D^{(u)}}^{1+\bar{A}:B'; \dots; B^{(u)}} \left(\begin{array}{c} y_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(1)}} (v_i - x_i)^{\beta_i'^{(1)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,1)}}} \right] \\ \vdots \\ y_u \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(u)}} (v_i - x_i)^{\beta_i'^{(u)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,u)}}} \right] \end{array} \right)$$

$$H \left(\begin{array}{c} z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i'^{(1)}} (v_i - x_i)^{\eta_i'} \rho_i^{(j,1)}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta_i'^{(r)}} (v_i - x_i)^{\eta_i^{(r)}} \rho_i^{(j,r)}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{array} \right)$$

$${}_p F_Q \left[(A_P); (B_Q); - \sum_{k=1}^l g_k \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\mu_i^{(k)}} (v_i - x_i)^{\theta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,k)}}} \right] \right] dx_1 \cdots dx_s$$

$$= \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{i=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\sum_{n'=0}^{\infty} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} B'_u(a)_{n'} \frac{B_p^{(\alpha, \beta; \tau, \mu)}(b + n', c - b)}{B(b, c - b)} \frac{z^{n'}}{n'!} y_1^{R_1} \cdots y_u^{R_u} \prod_{i=1}^s \left[(v_i - u_i)^{n'(a_i + b_i) + \sum_{k=1}^u R_k (\alpha_i^{(k)} + \beta_i^{(k)})} \right]$$

$$\prod_{i=1}^s \left[\prod_{j=1}^W \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)}} \prod_{j=W+1}^T \left(u_i U_i^{(j)} + V_i^{(j)} \right)^{-n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j,k)}} \right]$$

$$H_{\mathbf{p}+sT+P+2s, \mathbf{q}+sT+Q+s; |R':X}^{\mathbf{m}, \mathbf{n}+sT+P+2s; |R:Y} \left(\begin{array}{c|c} z_1 w_1 & \mathbb{A}, \mathbb{A}^* \\ \cdots & \cdot \\ \cdots & \cdot \\ z_r w_r & \cdot \\ g_1 W_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ g_l W_l & \cdot \\ G_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ G_T & \mathbb{B}, \mathbb{B}^* \end{array} \right) \quad (7.3)$$

under the same notations and conditions that (5.7)

and $B'_u = \frac{(-L)_{h_1 R_1 + \cdots + h_u R_u} B(E; R_1, \dots, R_u)}{R_1! \cdots R_u!}$; $B(L; R_1, \dots, R_u)$ is defined by (8.1)

Corollary 2

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[(x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$F_p^{(\alpha, \beta; \tau, \mu)} \left[a, b, c; z \prod_{i=1}^s \left[\frac{(x_i - u_i)^{a_i} (v_i - x_i)^{b_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{c_i^{(j)}}} \right] \right] R_n^{\alpha, \beta} \left[Z \prod_{j=1}^s \left[\frac{(x_i - u_i)^{\zeta_i} (v_i - x_i)^{\lambda_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\theta_i^{(j)}}} \right] \right]$$

$$F_{\bar{C}:D'; \dots; D^{(u)}}^{1+\bar{A}:B'; \dots; B^{(u)}} \left(\begin{array}{c} y_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(1)}} (v_i - x_i)^{\beta_i'^{(1)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,1)}}} \right] \\ \cdot \\ \cdot \\ y_u \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\alpha_i'^{(u)}} (v_i - x_i)^{\beta_i'^{(u)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,u)}}} \right] \end{array} \right)$$

$$\begin{aligned}
& H \left(\begin{array}{c} z_1 \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta'_i(1)} (v_i - x_i)^{\eta'_i} }{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[\frac{(x_i - u_i)^{\delta'_i(r)} (v_i - x_i)^{\eta_i^{(r)}} }{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{array} \right) dx_1 \cdots dx_s \\
&= \prod_{i=1}^s \left[(v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\rho_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\rho_i^{(j)}} \right] \sum_{n'=0}^{\infty} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \sum_{w, v, u, t, e, k_1, k_2} \\
& B_u(a)_{n'} \frac{B_p^{(\alpha, \beta; \tau, \mu)}(b + n', c - b)}{B(b, c - b)} \frac{z^{n'}}{n'!} y_1^{R_1} \cdots y_u^{R_u} \psi'(w, v, u, t, e, k_1, k_2) Z^R \prod_{i=1}^s \left[(v_i - u_i)^{n'(a_i + b_i) + \sum_{k=1}^u R_k (\alpha_i^{(k)} + \beta_i^{(k)})} \right] \\
& \prod_{i=1}^s \left[\prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{-n' c_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j, k)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{-n' e_i^{(j)} - \sum_{k=1}^u R_k \xi_i^{(j, k)}} \right] \\
& \prod_{j=1}^s \left[(v_i - u_i)^{n'(a_i + b_i) + \sum_{k=1}^u R_k (\alpha_i^{(k)} + \beta_i^{(k)})} \right] H_{\mathbf{p} + sT + 2s, \mathbf{q} + sT + s; |R': X}^{\mathbf{m}, \mathbf{n} + sT + 2s; |R: Y} \left(\begin{array}{c|c} z_1 w_1 & \mathbb{A}, A^* \\ \cdots & \cdot \\ \cdots & \cdot \\ z_r w_r & \cdot \\ \cdots & \cdot \\ G_1 & \cdot \\ \cdots & \cdot \\ \cdots & \cdot \\ G_T & \mathbb{B}, B^* \end{array} \right) \quad (7.4)
\end{aligned}$$

under the same notations and conditions that (6.7)

$$\text{and } B'_u = \frac{(-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u)}{R_1! \cdots R_u!}; \quad B(L; R_1, \dots, R_u) \text{ is defined by (8.1)}$$

8. Conclusion

In this paper we have evaluated two generalized multiple Eulerian integrals involving the modified multivariable H-functions defined by Prasad and Singh [5] with general arguments. The formulae established in this paper is very general nature. Thus, the results established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

- [1] Agrawal B.D. And Chaubey J.P. Certain derivation of generating relations for generalized polynomials. Indian J.

Pure and Appl. Math 10 (1980), page 1155-1157, ibid 11 (1981), page 357-359

[2] Fujiwara I. A unified presentation of classical orthogonal polynomials. Math. Japon. 11 (1966), page 133-148.

[3] Garg R. Unified multiple Eulerian integrals. Ganita Sandesh. Vol 15, no 2, 2001, page 2001.

[4] Goyal S.P. and Mathur T. On general multiple Eulerian Integrals and fractional integration. Vijnana. Parishad. Anusandhan. Patrika. Vol 46, no 3, 2003, page 231-245.

[5] Prasad Y.N. and Singh A.K. Basic properties of the transform involving and H-function of r-variables as kernel. Indian Acad Math, no 2, 1982, page 109-115.

[6] Raizada S.K. A study of unified representation of special functions of Mathematics Physics and their use in statistical and boundary value problem. Ph.D. Thesis, Bundelkhand University, Jhansi, India, 1991.

[7] Salim T.O. A serie formula of generalized class of polynomials associated with Laplace transform and fractional integral operators. J. Rajasthan. Acad. Phy. Sci. 1(3) (2002), page 167-176.

[8] Srivastava H.M. Agarwal P. and Jain S. Generating functions for the generalized Gauss hypergeometric functions. Applied Mathematics and computations 247 (2014), page 358-352

[9] Srivastava H.M. and Daoust M.C. Certain generalized Neumann expansions associated with Kampé de Fériet function. Nederl. Akad. Wetensch. Proc. Ser A72 = Indag Math 31(1969) page 449-457.

[10] Srivastava H.M. And Garg M. Some integral involving a general class of polynomials and multivariable H-function. Rev. Roumaine Phys. 32(1987), page 685-692.

[11] Srivastava H.M. Gupta K.C. And Goyal S.P. The H-functions of one and two variables with applications. South Asian Publishers Pvt Ltd 1982.

[12] Srivastava H.M. and Karlsson P.W. Multiple Gaussian Hypergeometric series. Ellis Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.

[13] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.