

## Eulerian integral associated with product of two multivariable I-functions and a class of polynomials of several variables

F.Y. AYANT<sup>1</sup>

<sup>1</sup> Teacher in High School , France  
 E-mail : fredericayant@gmail.com

**ABSTRACT**

The present paper is evaluated a new Eulerian integral associated with the product of two multivariable I-functions with general arguments and general class of polynomials. Several particular cases are given .

Keywords: Eulerian integral, multivariable I-function, generalized Lauricella function of several variables, multivariable H-function, generalized hypergeometric function, class of polynomials.

2010 Mathematics Subject Classification. 33C60, 82C31

### 1. Introduction

In this paper, we evaluate a new Eulerian integral of most general characters associated with the products of two multivariable I-functions defined by Prasad [1] and general class of polynomials of several variables, but of greater order. Several particular cases are given.

The multivariable I-function of r-variables is defined in term of multiple Mellin-Barnes type integral :

$$I(z_1, z_2, \dots, z_r) = I_{p_2, q_2, p_3, q_3; \dots; p_r, q_r; p^{(1)}, q^{(1)}; \dots; p^{(r)}, q^{(r)}}^{0, n_2; 0, n_3; \dots; 0, n_r; m^{(1)}, n^{(1)}; \dots; m^{(r)}, n^{(r)}} \left( \begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \middle| \begin{matrix} (a_{2j}; \alpha'_{2j}, \alpha''_{2j})_{1, p_2}; \dots; \\ \\ \\ (b_{2j}; \beta'_{2j}, \beta''_{2j})_{1, q_2}; \dots; \end{matrix} \right)$$

$$\left( \begin{matrix} (a_{rj}; \alpha_{rj}^{(1)}, \dots, \alpha_{rj}^{(r)})_{1, p_r}; (a_j^{(1)}, \alpha_j^{(1)})_{1, p^{(1)}}; \dots; (a_j^{(r)}, \alpha_j^{(r)})_{1, p^{(r)}} \\ \\ (b_{rj}; \beta_{rj}^{(1)}, \dots, \beta_{rj}^{(r)})_{1, q_r}; (b_j^{(1)}, \beta_j^{(1)})_{1, q^{(1)}}; \dots; (b_j^{(r)}, \beta_j^{(r)})_{1, q^{(r)}} \end{matrix} \right) \tag{1.1}$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \phi(s_1, \dots, s_r) \prod_{i=1}^r \theta_i(s_i) z_i^{s_i} ds_1 \dots ds_r \tag{1.2}$$

The defined integral of the above function, the existence and convergence conditions, see Y,N Prasad [1]. Throughout the present document, we assume that the existence and convergence conditions of the multivariable I-function.

The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|argz_i| < \frac{1}{2} \Omega_i \pi, \text{ where}$$

$$\Omega_i = \sum_{k=1}^{n^{(i)}} \alpha_k^{(i)} - \sum_{k=n^{(i)}+1}^{p^{(i)}} \alpha_k^{(i)} + \sum_{k=1}^{m^{(i)}} \beta_k^{(i)} - \sum_{k=m^{(i)}+1}^{q^{(i)}} \beta_k^{(i)} + \left( \sum_{k=1}^{n_2} \alpha_{2k}^{(i)} - \sum_{k=n_2+1}^{p_2} \alpha_{2k}^{(i)} \right) + \dots +$$

$$\left( \sum_{k=1}^{n_s} \alpha_{sk}^{(i)} - \sum_{k=n_s+1}^{p_s} \alpha_{sk}^{(i)} \right) - \left( \sum_{k=1}^{q_2} \beta_{2k}^{(i)} + \sum_{k=1}^{q_3} \beta_{3k}^{(i)} + \dots + \sum_{k=1}^{q_s} \beta_{sk}^{(i)} \right) \quad (1.3)$$

where  $i = 1, \dots, r$

The complex numbers  $z_i$  are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable I-function. We may establish the asymptotic expansion in the following convenient form :

$$I(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$I(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty$$

where  $k = 1, \dots, r : \alpha'_k = \min[\operatorname{Re}(b_j^{(k)} / \beta_j^{(k)})], j = 1, \dots, m_k$  and

$$\beta'_k = \max[\operatorname{Re}((a_j^{(k)} - 1) / \alpha_j^{(k)})], j = 1, \dots, n_k$$

We will use these following notations in this section :

$$U_r = p_2, q_2; p_3, q_3; \dots; p_{r-1}, q_{r-1}; V_r = 0, n_2; 0, n_3; \dots; 0, n_{r-1} \quad (1.4)$$

$$W_r = (p^{(1)}, q^{(1)}); \dots; (p^{(r)}, q^{(r)}); X_r = (m^{(1)}, n^{(1)}); \dots; (m^{(r)}, n^{(r)}) \quad (1.5)$$

$$A = (a_{2k}; \alpha_{2k}^{(1)}, \alpha_{2k}^{(2)}); \dots; (a_{(r-1)k}; \alpha_{(r-1)k}^{(1)}, \alpha_{(r-1)k}^{(2)}, \dots, \alpha_{(r-1)k}^{(r-1)}) \quad (1.6)$$

$$B = (b_{2k}; \beta_{2k}^{(1)}, \beta_{2k}^{(2)}); \dots; (b_{(r-1)k}; \beta_{(r-1)k}^{(1)}, \beta_{(r-1)k}^{(2)}, \dots, \beta_{(r-1)k}^{(r-1)}) \quad (1.7)$$

$$\mathfrak{A} = (a_{rk}; \alpha_{rk}^{(1)}, \alpha_{rk}^{(2)}, \dots, \alpha_{rk}^{(r)}) : \mathfrak{B} = (b_{rk}; \beta_{rk}^{(1)}, \beta_{rk}^{(2)}, \dots, \beta_{rk}^{(r)}) \quad (1.8)$$

$$A_1 = (a_k^{(1)}, \alpha_k^{(1)})_{1,p^{(1)}}; \dots; (a_k^{(r)}, \alpha_k^{(r)})_{1,p^{(r)}}; B_1 = (b_k^{(1)}, \beta_k^{(1)})_{1,p^{(1)}}; \dots; (b_k^{(r)}, \beta_k^{(r)})_{1,p^{(r)}} \quad (1.9)$$

The multivariable I-function of r-variables write :

$$I(z_1, \dots, z_r) = I_{U_r; 0; n_r; X_r}^{V_r; 0; n_r; X_r} \left( \begin{array}{c|c} z_1 & A; \mathfrak{A}; A_1 \\ \cdot & \\ \cdot & \\ \cdot & \\ \cdot & \\ z_r & B; \mathfrak{B}; B_1 \end{array} \right) \quad (1.10)$$

$$I(z'_1, \dots, z'_s) = I_{p'_2, q'_2, p'_3, q'_3; \dots; p'_s, q'_s; p'(1), q'(1); \dots; p'(s), q'(s)}^{0, n'_2; 0, n'_3; \dots; 0, n'_s; m'(1), n'(1); \dots; m'(s), n'(s)} \left( \begin{array}{c|l} z'_1 & (a'_{2j}; \alpha'_{2j}(1), \alpha'_{2j}(2))_{1, p'_2}; \dots; \\ \cdot & \\ \cdot & \\ \cdot & \\ z'_s & (b'_{2j}; \beta'_{2j}(1), \beta'_{2j}(2))_{1, q'_2}; \dots; \end{array} \right)$$

$$\left( \begin{array}{l} (a'_{sj}; \alpha'_{sj}(1), \dots, \alpha'_{sj}(s))_{1, p'_s} : (a'_j(1), \alpha'_j(1))_{1, p'(1)}; \dots; (a'_j(s), \alpha'_j(s))_{1, p'(s)} \\ (b'_{sj}; \beta'_{sj}(1), \dots, \beta'_{sj}(s))_{1, q'_s} : (b'_j(1), \beta'_j(1))_{1, q'(1)}; \dots; (b'_j(s), \beta'_j(s))_{1, q'(s)} \end{array} \right) \quad (1.11)$$

$$= \frac{1}{(2\pi\omega)^s} \int_{L_1} \dots \int_{L_s} \psi(t_1, \dots, t_s) \prod_{i=1}^s \xi_i(t_i) z_i^{t_i} dt_1 \dots dt_s \quad (1.12)$$

The defined integral of the above function, the existence and convergence conditions, see Y,N Prasad [1]. Throughout the present document, we assume that the existence and convergence conditions of the multivariable I-function.

The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$\begin{aligned} \text{where } |arg z'_i| < \frac{1}{2} \Omega'_i \pi, \\ \Omega'_i = \sum_{k=1}^{n'(i)} \alpha'_k(i) - \sum_{k=n'(i)+1}^{p'(i)} \alpha'_k(i) + \sum_{k=1}^{m'(i)} \beta'_k(i) - \sum_{k=m'(i)+1}^{q'(i)} \beta'_k(i) + \left( \sum_{k=1}^{n'_2} \alpha'_{2k}(i) - \sum_{k=n_2+1}^{p'_2} \alpha'_{2k}(i) \right) \\ + \dots + \left( \sum_{k=1}^{n'_s} \alpha'_{sk}(i) - \sum_{k=n'_s+1}^{p'_s} \alpha'_{sk}(i) \right) - \left( \sum_{k=1}^{q'_2} \beta'_{2k}(i) + \sum_{k=1}^{q'_3} \beta'_{3k}(i) + \dots + \sum_{k=1}^{q'_s} \beta'_{sk}(i) \right) \end{aligned} \quad (1.13)$$

where  $i = 1, \dots, s$

The complex numbers  $z_i$  are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable I-function.

We may establish the the asymptotic expansion in the following convenient form :

$$I(z'_1, \dots, z'_s) = O(|z'_1|^{\alpha'_1}, \dots, |z'_s|^{\alpha'_s}), \max(|z'_1|, \dots, |z'_s|) \rightarrow 0$$

$$I(z'_1, \dots, z'_s) = O(|z'_1|^{\beta'_1}, \dots, |z'_s|^{\beta'_s}), \min(|z'_1|, \dots, |z'_s|) \rightarrow \infty$$

where  $k = 1, \dots, s : \alpha''_k = \min[Re(b_j^{(k)}/\beta_j^{(k)})], j = 1, \dots, m'_k$  and

$$\beta''_k = \max[Re((a_j^{(k)} - 1)/\alpha_j^{(k)})], j = 1, \dots, n'_k$$

We will use these following notations in this section :

$$U_s = p'_2, q'_2; p'_3, q'_3; \dots; p'_{s-1}, q'_{s-1}; V_s = 0, n_2; 0, n_3; \dots; 0, n_{s-1} \quad (1.14)$$

$$W_s = (p'^{(1)}, q'^{(1)}); \dots; (p'^{(s)}, q'^{(s)}); X_s = (m'^{(1)}, n'^{(1)}); \dots; (m'^{(s)}, n'^{(s)}) \quad (1.15)$$

$$A' = (a'_{2k}; \alpha'^{(1)}_{2k}, \alpha'^{(2)}_{2k}); \dots; (a'_{(s-1)k}; \alpha'^{(1)}_{(s-1)k}, \alpha'^{(2)}_{(s-1)k}, \dots, \alpha'^{(s-1)}_{(s-1)k}) \quad (1.16)$$

$$B' = (b'_{2k}; \beta'^{(1)}_{2k}, \beta'^{(2)}_{2k}); \dots; (b'_{(s-1)k}; \beta'^{(1)}_{(s-1)k}, \beta'^{(2)}_{(s-1)k}, \dots, \beta'^{(s-1)}_{(s-1)k}) \quad (1.17)$$

$$\mathfrak{A}' = (a'_{sk}; \alpha'^{(1)}_{sk}, \alpha'^{(2)}_{sk}, \dots, \alpha'^{(s)}_{sk}) : \mathfrak{B}' = (b'_{sk}; \beta'^{(1)}_{sk}, \beta'^{(2)}_{sk}, \dots, \beta'^{(s)}_{sk}) \quad (1.18)$$

$$A'_1 = (a'_k{}^{(1)}, \alpha'_k{}^{(1)})_{1,p'(1)}; \dots; (a'_k{}^{(s)}, \alpha'_k{}^{(s)})_{1,p'(s)}; B'_1 = (b'_k{}^{(1)}, \beta'_k{}^{(1)})_{1,p'(1)}; \dots; (b'_k{}^{(s)}, \beta'_k{}^{(s)})_{1,p'(s)} \quad (1.19)$$

The multivariable I-function write :

$$I(z'_1, \dots, z'_s) = I_{U_s; p'_s, q'_s; W_s}^{V_s; 0, n'_s; X_s} \left( \begin{array}{c} z'_1 \\ \vdots \\ \vdots \\ z'_s \end{array} \middle| \begin{array}{l} A'; \mathfrak{A}'; A'_1 \\ \\ B'; \mathfrak{B}'; B'_1 \end{array} \right) \quad (1.20)$$

Srivastava and Garg introduced and defined a general class of multivariable polynomials [3] as follows

$$S_L^{h_1, \dots, h_u} [z_1, \dots, z_u] = \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{z_1^{R_1} \dots z_u^{R_u}}{R_1! \dots R_u!} \quad (1.21)$$

## 2. Integral representation of generalized Lauricella function of several variables

The following generalized hypergeometric function in terms of multiple contour integrals is also required [4 ,page 39 eq .30]

$$\begin{aligned} & \frac{\prod_{j=1}^P \Gamma(A_j)}{\prod_{j=1}^Q \Gamma(B_j)} {}_P F_Q [(A_P); (B_Q); -(x_1 + \dots + x_r)] \\ &= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \frac{\prod_{j=1}^P \Gamma(A_j + s_1 + \dots + s_r)}{\prod_{j=1}^Q \Gamma(B_j + s_1 + \dots + s_r)} \Gamma(-s_1) \dots \Gamma(-s_r) x_1^{s_1} \dots x_r^{s_r} ds_1 \dots ds_r \end{aligned} \quad (2.1)$$

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of  $\Gamma(A_j + s_1 + \dots + s_r)$  are separated from those of  $\Gamma(-s_j), j = 1, \dots, r$ . The above result (1.23) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of  $\Gamma(-s_j), j = 1, \dots, r$

The Lauricella function  $F_D^{(k)}$  is defined as

$$F_D^{(k)} [a, b_1, \dots, b_k; c; x_1, \dots, x_k] = \frac{\Gamma(c)}{\Gamma(a) \prod_{j=1}^k \Gamma(b_j)} \frac{1}{(2\pi\omega)^k} \int_{L_1} \dots \int_{L_k} \frac{\Gamma\left(a + \sum_{j=1}^k \zeta_j\right) \Gamma(b_1 + \zeta_1), \dots, \Gamma(b_k + \zeta_k)}{\Gamma\left(c + \sum_{j=1}^k \zeta_j\right)} \prod_{j=1}^k \Gamma(-\zeta_j) (-x_j)^{\zeta_j} d\zeta_1 \dots d\zeta_k \quad (2.2)$$

where  $\max[|\arg(-x_1)|, \dots, |\arg(-x_k)|] < \pi, c \neq 0, -1, -2, \dots$

In order to evaluate a number of integrals of multivariable I-function, we first establish the formula

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j} dt = (b-a)^{\alpha+\beta-1} B(\alpha, \beta) \prod_{j=1}^k (af_j + g_j)^{\sigma_j} \times F_D^{(k)} \left[ \alpha, -\sigma_1, \dots, -\sigma_k; \alpha + \beta; -\frac{(b-a)f_1}{af_1 + g_1}, \dots, -\frac{(b-a)f_k}{af_k + g_k} \right] \quad (2.3)$$

where  $a, b \in \mathbb{R} (a < b), \alpha, \beta, f_i, g_i, \sigma_i \in \mathbb{C}, (i = 1, \dots, k); \min(\operatorname{Re}(\alpha), \operatorname{Re}(\beta)) > 0$  and

$$\max_{1 \leq j \leq k} \left\{ \left| \frac{(b-a)f_i}{af_i + g_i} \right| \right\} < 1$$

$F_D^{(k)}$  is a Lauricella's function of  $k$ -variables, see Srivastava et al ([5], page60)

The formula (2.2) can be establish by expanding  $\prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$  by means of the formula :

$$(1-z)^{-\alpha} = \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r!} z^r (|z| < 1) \quad (2.4)$$

integrating term by term with the help of the integral given by Saigo and Saxena [2, page 93, eq.(3.2)] and applying the definition of the Lauricella function  $F_D^{(k)}$  [4, page 454].

### 3. Eulerian integral

In this section , we note :

$$U = p_2, q_2; p_3, q_3; \dots; p_{r-1}, q_{r-1}; p'_2, q'_2; p'_3, q'_3; \dots; p'_{s-1}, q'_{s-1}; 0, 0; \dots; 0, 0; 0, 0; \dots; 0, 0 \quad (3.1)$$

$$V = 0, n_2; 0, n_3; \dots; 0, n_{r-1}; 0, n'_2; 0, n'_3; \dots; 0, n'_{s-1}; 0, 0; \dots; 0, 0; 0, 0; \dots; 0, 0 \quad (3.2)$$

$$X = m^{(1)}, n^{(1)}; \dots; m^{(r)}, n^{(r)}; m'^{(1)}, n'^{(1)}; \dots; m'^{(s)}, n'^{(s)}; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0 \quad (3.3)$$

$$Y = p^{(1)}, q^{(1)}; \dots; p^{(r)}, q^{(r)}; p'^{(1)}, q'^{(1)}; \dots; p'^{(s)}, q'^{(s)}; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1 \quad (3.4)$$

$$A = (a_{2k}; \alpha_{2k}^{(1)}, \alpha_{2k}^{(2)}); \dots; (a_{(r-1)k}; \alpha_{(r-1)k}^{(1)}, \alpha_{(r-1)k}^{(2)}, \dots, \alpha_{(r-1)k}^{(r-1)}); (a'_{2k}; \alpha'_{2k}^{(1)}, \alpha'_{2k}^{(2)}); \dots;$$

$$(a'_{(s-1)k}; \alpha'_{(s-1)k}{}^{(1)}, \alpha'_{(s-1)k}{}^{(2)}, \dots, \alpha'_{(s-1)k}{}^{(s-1)}) \quad (3.5)$$

$$B = (b_{2k}; \beta_{2k}^{(1)}, \beta_{2k}^{(2)}); \dots; (b_{(r-1)k}; \beta_{(r-1)k}^{(1)}, \beta_{(r-1)k}^{(2)}, \dots, \beta_{(r-1)k}^{(r-1)}); (b'_{2k}; \beta'_{2k}{}^{(1)}, \beta'_{2k}{}^{(2)}); \dots;$$

$$(b'_{(s-1)k}; \beta'_{(s-1)k}{}^{(1)}, \beta'_{(s-1)k}{}^{(2)}, \dots, \beta'_{(s-1)k}{}^{(s-1)}) \quad (3.6)$$

$$\mathfrak{A} = (a_{rk}; \alpha_{rk}^{(1)}, \alpha_{rk}^{(2)}, \dots, \alpha_{rk}^{(r)}, 0, \dots, 0, 0, \dots, 0, 0, \dots, 0) \quad (3.7)$$

$$\mathfrak{A}' = (a'_{sk}; 0, \dots, 0, \alpha'_{sk}{}^{(1)}, \alpha'_{sk}{}^{(2)}, \dots, \alpha'_{sk}{}^{(s)}, 0, \dots, 0, 0, \dots, 0) \quad (3.8)$$

$$\mathfrak{B} = (b_{rk}; \beta_{rk}^{(1)}, \beta_{rk}^{(2)}, \dots, \beta_{rk}^{(r)}, 0, \dots, 0, 0, \dots, 0, 0, \dots, 0) \quad (3.9)$$

$$\mathfrak{B}' = (b'_{sk}; 0, \dots, 0, \beta'_{sk}{}^{(1)}, \beta'_{sk}{}^{(2)}, \dots, \beta'_{sk}{}^{(s)}, 0, \dots, 0, 0, \dots, 0) \quad (3.10)$$

$$\mathfrak{A}_1 = (a_k^{(1)}, \alpha_k^{(1)})_{1,p^{(1)}}; \dots; (a_k^{(r)}, \alpha_k^{(r)})_{1,p^{(r)}}; (a'_k{}^{(1)}, \alpha'_k{}^{(1)})_{1,p^{(1)}}; \dots; (a'_k{}^{(s)}, \alpha'_k{}^{(s)})_{1,p^{(s)}};$$

$$(1, 0); \dots; (1, 0); (1, 0); \dots; (1, 0) \quad (3.11)$$

$$\mathfrak{B}_1 = (b_k^{(1)}, \beta_k^{(1)})_{1,p^{(1)}}; \dots; (b_k^{(r)}, \beta_k^{(r)})_{1,p^{(r)}}; (b'_k{}^{(1)}, \beta'_k{}^{(1)})_{1,p^{(1)}}; \dots; (b'_k{}^{(s)}, \beta'_k{}^{(s)})_{1,p^{(s)}};$$

$$(0, 1); \dots; (0, 1); (0, 1); \dots; (0, 1) \quad (3.12)$$

$$K_1 = (1 - \alpha - \sum_{i=1}^u R_i(\mu_i + \mu'_i); \mu_1, \dots, \mu_r, \mu'_1, \dots, \mu'_s, 1, \dots, 1, v_1, \dots, v_l) \quad (3.13)$$

$$K_2 = (1 - \beta - \sum_{i=1}^u R_i(\rho_i + \rho'_i); \rho_1, \dots, \rho_r, \rho'_1, \dots, \rho'_s, 0, \dots, 0, \tau_1, \dots, \tau_l) \quad (3.14)$$

$$K_P = [1 - A_j; 0, \dots, 0, 0, \dots, 0, 0, \dots, 0, 1, \dots, 1]_{1,P} \quad (3.15)$$

$$K_j = [1 + \sigma_j - \sum_{i=1}^u R_i(\lambda_i^{(j)} + \lambda'_i{}^{(j)}); \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j^{(1)}, \dots, \lambda_j^{(s)}, 0, \dots, 1, \dots, 0, \zeta'_j, \dots, \zeta_j^{(l)}]_{1,k} \quad (3.16)$$

j

$$L_1 = (1 - \alpha - \beta - \sum_{i=1}^u R_i(\mu_i + \mu'_i + \rho_i + \rho'_i); \mu_1 + \rho_1, \dots, \mu_r + \rho_r, \mu'_1 + \rho'_1, \dots, \mu'_s + \rho'_s, 1, \dots, 1, v_1 + \tau_1, \dots, v_l + \tau_l) \quad (3.17)$$

$$L_Q = [1 - B_j; 0, \dots, 0, 0, \dots, 0, 0, \dots, 0, 1, \dots, 1]_{1,Q} \quad (3.18)$$

$$L_j = [1 + \sigma_j - \sum_{i=1}^u R_i(\lambda_i^{(j)} + \lambda_i'^{(j)}); \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j^{(1)}, \dots, \lambda_j^{(s)}, 0, \dots, 0, \zeta_j', \dots, \zeta_j^{(l)}]_{1,k} \quad (3.19)$$

We have the following result

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$S_L^{h_1, \dots, h_u} \left( \begin{array}{c} x_1 (t-a)^{\mu_1 + \mu'_1} (b-t)^{\rho_1 + \rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)} - \lambda_j'^{(1)}} \\ \vdots \\ x_u (t-a)^{\mu_u + \mu'_u} (b-t)^{\rho_u + \rho'_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)} - \lambda_j'^{(u)}} \end{array} \right)$$

$$I_{U_r: p_r, q_r; W_r}^{V_r; 0, n_r; X_r} \left( \begin{array}{c} z_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{array} \right)$$

$$I_{U_s: p'_s, q'_s; W_s}^{V_s; 0, n'_s; X_s} \left( \begin{array}{c} z'_1 (t-a)^{\mu'_1} (b-t)^{\rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(1)}} \\ \vdots \\ z'_s (t-a)^{\mu'_s} (b-t)^{\rho'_s} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(s)}} \end{array} \right)$$

$${}_P F_Q \left[ (A_P); (B_Q); - \sum_{i=1}^l z_i'' (t-a)^{\nu_i} (b-t)^{\tau_i} \prod_{j=1}^k (f_j t + g_j)^{-\zeta_j^{(i)}} \right] dt =$$

$$(b-a)^{\alpha + \beta + \sum_{i=1}^u R_i(\mu_i + \mu'_i + \rho_i + \rho'_i)} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^k (a f_j + g_j)^{\sigma_j - \sum_{i=1}^u R_i(\lambda_j^{(i)} + \lambda_j'^{(i)})}$$

$$\sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{x_1^{R_1} \dots x_u^{R_u}}{R_1! \dots R_u!}$$



$$\begin{aligned}
(\mathbf{E}) \Omega_i &= \sum_{k=1}^{n^{(i)}} \alpha_k^{(i)} - \sum_{k=n^{(i)}+1}^{p^{(i)}} \alpha_k^{(i)} + \sum_{k=1}^{m^{(i)}} \beta_k^{(i)} - \sum_{k=m^{(i)}+1}^{q^{(i)}} \beta_k^{(i)} + \left( \sum_{k=1}^{n_2} \alpha_{2k}^{(i)} - \sum_{k=n_2+1}^{p_2} \alpha_{2k}^{(i)} \right) + \dots + \\
&\left( \sum_{k=1}^{n_s} \alpha_{sk}^{(i)} - \sum_{k=n_s+1}^{p_s} \alpha_{sk}^{(i)} \right) - \left( \sum_{k=1}^{q_2} \beta_{2k}^{(i)} + \sum_{k=1}^{q_3} \beta_{3k}^{(i)} + \dots + \sum_{k=1}^{q_s} \beta_{sk}^{(i)} \right) - \mu_i - \rho_i \\
&- \sum_{j=1}^k \lambda_j^{(i)} > 0 \quad (i = 1, \dots, r)
\end{aligned}$$

$$\begin{aligned}
\Omega'_i &= \sum_{k=1}^{n'^{(i)}} \alpha'_k{}^{(i)} - \sum_{k=n'^{(i)}+1}^{p'^{(i)}} \alpha'_k{}^{(i)} + \sum_{k=1}^{m'^{(i)}} \beta'_k{}^{(i)} - \sum_{k=m'^{(i)}+1}^{q'^{(i)}} \beta'_k{}^{(i)} + \left( \sum_{k=1}^{n'_2} \alpha'_{2k}{}^{(i)} - \sum_{k=n'_2+1}^{p'_2} \alpha'_{2k}{}^{(i)} \right) + \\
&\dots + \left( \sum_{k=1}^{n'_s} \alpha'_{sk}{}^{(i)} - \sum_{k=n'_s+1}^{p'_s} \alpha'_{sk}{}^{(i)} \right) - \left( \sum_{k=1}^{q'_2} \beta'_{2k}{}^{(i)} + \sum_{k=1}^{q'_3} \beta'_{3k}{}^{(i)} + \dots + \sum_{k=1}^{q'_s} \beta'_{sk}{}^{(i)} \right) - \mu'_i - \rho'_i \\
&- \sum_{j=1}^k \lambda'_j{}^{(i)} > 0 \quad (i = 1, \dots, s)
\end{aligned}$$

$$(\mathbf{F}) \left| \arg \left( z_i \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right) \right| < \frac{1}{2} \Omega_i \pi \quad (a \leq t \leq b; i = 1, \dots, r)$$

$$\left| \arg \left( z_i \prod_{j=1}^k (f_j t + g_j)^{-\lambda'_j{}^{(i)}} \right) \right| < \frac{1}{2} \Omega_i \pi \quad (a \leq t \leq b; i = 1, \dots, s)$$

(G)  $P \leq Q + 1$ . The equality holds, when , in addition,

$$\text{either } P > Q \text{ and } \left| z_i \left( \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right) \right|^{\frac{1}{Q-P}} < 1 \quad (a \leq t \leq b)$$

$$\text{or } P \leq Q \text{ and } \max_{1 \leq i \leq k} \left[ \left| \left( z'_i \prod_{j=1}^k (f_j t + g_j)^{-\lambda'_j{}^{(i)}} \right) \right| \right] < 1 \quad (a \leq t \leq b)$$

### Proof

First expressing the class of polynomial in serie with the help of (1.21), expressing the I-function of r-variables by the Mellin-Barnes contour integral with the help of the equation (1.12), the generalized hypergeometric function  ${}_P F_Q(\cdot)$  in Mellin-Barnes contour integral with the help of (2.1). Now collect the power of  $(f_j t + g_j)$  with  $j = 1, \dots, k$  and use

the equations (2.1) and (2.2) and we obtain  $k$ -Mellin-Barnes contour integral. Changing the order of integration and summation (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process) and interpreting  $(r + s + k + l)$ -Mellin-barnes contour integral in multivariable I-function of Prasad, we obtain the desired result.

#### 4. Particular cases

$$\mathbf{a)} \text{ If } B(L; R_1, \dots, R_u) = \frac{\prod_{j=1}^A (a_j)_{R_1 \theta'_j + \dots + R_u \theta_j^{(u)}} \prod_{j=1}^{B'} (b'_j)_{R_1 \phi'_j} \dots \prod_{j=1}^{B^{(u)}} (b_j^{(u)})_{R_u \phi_j^{(u)}}}{\prod_{j=1}^C (c_j)_{m_1 \psi'_j + \dots + m_u \psi_j^{(u)}} \prod_{j=1}^{D'} (d'_j)_{R_1 \delta'_j} \dots \prod_{j=1}^{D^{(u)}} (d_j^{(u)})_{R_u \delta_j^{(u)}}} \quad (4.1)$$

then the general class of multivariable polynomial  $S_L^{R_1, \dots, R_u} [x_1, \dots, x_u]$  reduces to generalized Lauricella function defined by Srivastava et al [6].

$$F_{C:D'; \dots; D^{(s)}}^{1+A:B'; \dots; B^{(s)}} \left( \begin{matrix} x_1 \\ \dots \\ \dots \\ x_u \end{matrix} \middle| \begin{matrix} [(-L); R_1, \dots, R_u] [(a); \theta', \dots, \theta^{(u)}] : [(b'); \phi']; \dots; [(b^{(u)}), \phi^{(u)}] \\ [(c); \psi', \dots, \psi^{(u)}] : [(d'); \delta']; \dots; [(d^{(u)}), \delta^{(u)}] \end{matrix} \right) \quad (4.2)$$

and we have the formula

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$F_{C:D'; \dots; D^{(t)}}^{1+A:B'; \dots; B^{(t)}} \left( \begin{matrix} x_1 (t-a)^{\mu_1 + \mu'_1} (b-t)^{\rho_1 + \rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)} - \lambda_j'^{(1)}} \\ \vdots \\ x_u (t-a)^{\mu_u + \mu'_u} (b-t)^{\rho_u + \rho'_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)} - \lambda_j'^{(u)}} \end{matrix} \right)$$

$$\left( \begin{matrix} [(-L); R_1, \dots, R_u] [(a); \theta', \dots, \theta^{(u)}] : [(b'); \phi']; \dots; [(b^{(u)}), \phi^{(u)}] \\ [(c); \psi', \dots, \psi^{(u)}] : [(d'); \delta']; \dots; [(d^{(u)}), \delta^{(u)}] \end{matrix} \right)$$

$$I_{U_r: p_r, q_r; W_r}^{V_r; 0, n_r; X_r} \left( \begin{matrix} z_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$$I_{U_s; p'_s, q'_s; W_s}^{V_s; 0, n'_s; X_s} \left( \begin{array}{c} z'_1(t-a)^{\mu'_1} (b-t)^{\rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda'_j(1)} \\ \vdots \\ z'_s(t-a)^{\mu'_s} (b-t)^{\rho'_s} \prod_{j=1}^k (f_j t + g_j)^{-\lambda'_j(s)} \end{array} \right)$$

$${}_P F_Q \left[ (A_P); (B_Q); - \sum_{i=1}^l z''_i (t-a)^{v_i} (b-t)^{\tau_i} \prod_{j=1}^k (f_j t + g_j)^{-\zeta_j^{(i)}} \right] dt =$$

$$(b-a)^{\alpha+\beta+\sum_1^u R_i(\mu_i+\mu'_i+\rho_i+\rho'_i)} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^k (af_j + g_j)^{\sigma_j + \sum_{i=1}^u R_i(\lambda_j^{(i)} + \lambda'_j{}^{(i)})}$$

$$\sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{x_1^{R_1} \dots x_u^{R_u}}{R_1! \dots R_u!}$$

$$I_{U: p_r + p'_s + l + k + 2, q_r + q'_s + l + k + 1; Y}^{V: 0, n_r + n'_s + l + k + 2; X} \left( \begin{array}{c} \frac{z_1(b-a)^{\mu_1+\rho_1}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(1)}}} \\ \vdots \\ \frac{z_r(b-a)^{\mu_r+\rho_r}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(r)}}} \\ \frac{z'_1(b-a)^{\mu'_1+\rho'_1}}{\prod_{j=1}^k (af_j + g_j)^{\lambda'_j(1)}} \\ \vdots \\ \frac{z'_s(b-a)^{\mu'_s+\rho'_s}}{\prod_{j=1}^k (af_j + g_j)^{\lambda'_j(s)}} \\ \frac{(b-a)f_1}{af_1 + g_1} \\ \vdots \\ \frac{(b-a)f_k}{af_k + g_k} \\ \frac{z''_1(b-a)^{\tau_1+v_1}}{\prod_{j=1}^k (af_j + g_j)^{\zeta_j^{(1)}}} \\ \vdots \\ \frac{z''_l(b-a)^{\tau_l+v_l}}{\prod_{j=1}^k (af_j + g_j)^{\zeta_j^{(l)}}} \end{array} \middle| \begin{array}{l} A ; K_1, K_2, K_P, K_j, \mathfrak{A}, \mathfrak{A}'; A' \\ \vdots \\ B ; L_1, L_j, L_Q, \mathfrak{B}, \mathfrak{B}'; B' \end{array} \right) \quad (4.3)$$

$$\text{where } B(L; R_1, \dots, R_u) = \frac{\prod_{j=1}^A (a_j)_{R_1 \theta'_j + \dots + R_u \theta_j^{(u)}} \prod_{j=1}^{B'} (b'_j)_{R_1 \phi'_j} \dots \prod_{j=1}^{B^{(u)}} (b_j^{(u)})_{R_u \phi_j^{(u)}}}{\prod_{j=1}^C (c_j)_{m_1 \psi'_j + \dots + m_u \psi_j^{(u)}} \prod_{j=1}^{D'} (d'_j)_{R_1 \delta'_j} \dots \prod_{j=1}^{D^{(u)}} (d_j^{(u)})_{R_u \delta_j^{(u)}}} \quad (4.4)$$

b) If  $U_r = V_r = U_s = V_s = U = V = A = B = 0$ , the multivariable I-function defined by Prasad degenerate in

**multivariable H-function defined by Srivastava et al [6].**

We have the following result (H-function of  $r + s + k + l$  variables).

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$S_L^{h_1, \dots, h_u} \left( \begin{matrix} x_1 (t-a)^{\mu_1 + \mu'_1} (b-t)^{\rho_1 + \rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)} - \lambda_j'^{(1)}} \\ \vdots \\ x_u (t-a)^{\mu_u + \mu'_u} (b-t)^{\rho_u + \rho'_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)} - \lambda_j'^{(u)}} \end{matrix} \right)$$

$$H_{p_r, q_r; W_r}^{0, n_r; X_r} \left( \begin{matrix} z_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$$H_{p'_s, q'_s; W_s}^{0, n'_s; X_s} \left( \begin{matrix} z'_1 (t-a)^{\mu'_1} (b-t)^{\rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(1)}} \\ \vdots \\ z'_s (t-a)^{\mu'_s} (b-t)^{\rho'_s} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(s)}} \end{matrix} \right)$$

$${}_P F_Q \left[ (A_P); (B_Q); - \sum_{i=1}^l z_i'' (t-a)^{v_i} (b-t)^{\tau_i} \prod_{j=1}^k (f_j t + g_j)^{-\zeta_j^{(i)}} \right] dt =$$

$$(b-a)^{\alpha + \beta + \sum_{i=1}^l R_i (\mu_i + \mu'_i + \rho_i + \rho'_i)} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^k (a f_j + g_j)^{\sigma_j + \sum_{i=1}^l R_i (\lambda_j^{(i)} + \lambda_j'^{(i)})}$$

$$\sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{x_1^{R_1} \dots x_u^{R_u}}{R_1! \dots R_u!}$$

$$H_{p_r+n_r+n'_s+l+k+2; X}^{0, n_r, n'_s+l+k+2, q_r+q'_s+l+k+1; Y} \left( \begin{array}{c} \frac{z_1(b-a)^{\mu_1+\rho_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(1)}}} \\ \dots \\ \frac{z_r(b-a)^{\mu_r+\rho_r}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(r)}}} \\ \frac{z'_1(b-a)^{\mu'_1+\rho'_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda'_j^{(1)}}} \\ \dots \\ \frac{z'_s(b-a)^{\mu'_s+\rho'_s}}{\prod_{j=1}^k (af_j+g_j)^{\lambda'_j^{(s)}}} \\ \frac{(b-a)f_1}{af_1+g_1} \\ \dots \\ \frac{(b-a)f_k}{af_k+g_k} \\ \frac{z''_1(b-a)^{\tau_1+\nu_1}}{\prod_{j=1}^k (af_j+g_j)^{\zeta_j^{(1)}}} \\ \dots \\ \frac{z''_l(b-a)^{\tau_l+\nu_l}}{\prod_{j=1}^k (af_j+g_j)^{\zeta_j^{(l)}}} \end{array} \middle| \begin{array}{c} K_1, K_2, K_P, K_j, \mathfrak{A}, \mathfrak{A}'; A' \\ \dots \\ L_1, L_j, L_Q, \mathfrak{B}, \mathfrak{B}'; B' \end{array} \right) \quad (4.5)$$

under the same conditions (3.19) where  $U_r = V_r = U_s = V_s = U = V = A = B = 0$

c) d) Let  $z'_1, \dots, z'_z \rightarrow 0$  and  $\lambda_j^{(i)} \rightarrow 0, (i = 1, \dots, s; j = 1, \dots, k)$  we obtain I-function of  $(r+k+l)$ - variables. We note

In this section, we denote :

$$U = p_2, q_2; p_3, q_3; \dots; p_{r-1}, q_{r-1}; 0, 0; 0, 0; \dots; 0, 0; 0, 0; \dots; 0, 0 \quad (4.6)$$

$$V = 0, n_2; 0, n_3; \dots; 0, n_{r-1}; 0, 0; \dots; 0, 0; 0, 0; \dots; 0, 0 \quad (4.7)$$

$$X = m^{(1)}, n^{(1)}; \dots; m^{(r)}, n^{(r)}; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0 \quad (4.8)$$

$$Y = p^{(1)}, q^{(1)}; \dots; p^{(r)}, q^{(r)}; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1 \quad (4.9)$$

$$A = (a_{2k}; \alpha_{2k}^{(1)}, \alpha_{2k}^{(2)}); \dots; (a_{(r-1)k}; \alpha_{(r-1)k}^{(1)}, \alpha_{(r-1)k}^{(2)}, \dots, \alpha_{(r-1)k}^{(r-1)}) \quad (4.10)$$

$$B = (b_{2k}; \beta_{2k}^{(1)}, \beta_{2k}^{(2)}); \dots; (b_{(r-1)k}; \beta_{(r-1)k}^{(1)}, \beta_{(r-1)k}^{(2)}, \dots, \beta_{(r-1)k}^{(r-1)}) \quad (4.11)$$

$$\mathfrak{A} = (a_{rk}; \alpha_{rk}^{(1)}, \alpha_{rk}^{(2)}, \dots, \alpha_{rk}^{(r)}, 0, \dots, 0, 0 \dots, 0) \quad (4.12)$$

$$\mathfrak{B} = (b_{rk}; \beta_{rk}^{(1)}, \beta_{rk}^{(2)}, \dots, \beta_{rk}^{(r)}, 0, \dots, 0, 0, \dots, 0) \quad (4.13)$$

$$\mathfrak{A}_1 = (a_k^{(1)}, \alpha_k^{(1)})_{1,p^{(1)}}; \cdots; (a_k^{(r)}, \alpha_k^{(r)})_{1,p^{(r)}}; (1, 0); \cdots; (1, 0); (1, 0); \cdots; (1, 0) \quad (4.14)$$

$$\mathfrak{B}_1 = (b_k^{(1)}, \beta_k^{(1)})_{1,p^{(1)}}; \cdots; (b_k^{(r)}, \beta_k^{(r)})_{1,p^{(r)}}; (0, 1); \cdots; (0, 1); (0, 1); \cdots; (0, 1) \quad (4.15)$$

$$K_1 = (1 - \alpha - \sum_{i=1}^u R_i \mu_i; \mu_1, \cdots, \mu_r; 1, \cdots, 1, v_1, \cdots, v_l) \quad (4.16)$$

$$K_2 = (1 - \beta - \sum_{i=1}^u R_i \rho_i; \rho_1, \cdots, \rho_r; 0, \cdots, 0, \tau_1, \cdots, \tau_l) \quad (4.17)$$

$$K_3 = [1 - A_j; 0, \cdots, 0, 0, \cdots, 0, 1, \cdots, 1]_{1,P} \quad (4.18)$$

$$K_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_i^{(j)}; \lambda_j^{(1)}, \cdots, \lambda_j^{(r)}; 0, \cdots, 1, \cdots, 0, \zeta'_j, \cdots, \zeta_j^{(l)}]_{1,k} \quad (4.19)$$

j

$$L_1 = (1 - \alpha - \beta - \sum_{i=1}^u R_i (\mu_i + \rho_i); \mu_1 + \rho_1, \cdots, \mu_r + \rho_r; 1, \cdots, 1, v_1 + \tau_1, \cdots, v_l + \tau_l) \quad (4.20)$$

$$L_2 = [1 - B_j; 0, \cdots, 0, 0, \cdots, 0, 1, \cdots, 1]_{1,Q} \quad (4.21)$$

$$L_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_i^{(j)}; \lambda_j^{(1)}, \cdots, \lambda_j^{(r)}; 0, \cdots, 0, \zeta'_j, \cdots, \zeta_j^{(l)}]_{1,k} \quad (4.22)$$

We have the following integral :

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$S_L^{h_1, \dots, h_u} \left( \begin{array}{c} x_1 (t-a)^{\mu_1 + \mu'_1} (b-t)^{\rho_1 + \rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ x_u (t-a)^{\mu_u + \mu'_u} (b-t)^{\rho_u + \rho'_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)}} \end{array} \right)$$

$$I_{U_r; p_r, q_r; W_r}^{V_r; 0, n_r; X_r} \left( \begin{array}{c} z_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{array} \right)$$



$$\begin{aligned}
\text{(E)} \quad \Omega_i &= \sum_{k=1}^{n^{(i)}} \alpha_k^{(i)} - \sum_{k=n^{(i)}+1}^{p^{(i)}} \alpha_k^{(i)} + \sum_{k=1}^{m^{(i)}} \beta_k^{(i)} - \sum_{k=m^{(i)}+1}^{q^{(i)}} \beta_k^{(i)} + \left( \sum_{k=1}^{n_2} \alpha_{2k}^{(i)} - \sum_{k=n_2+1}^{p_2} \alpha_{2k}^{(i)} \right) + \dots + \\
&\left( \sum_{k=1}^{n_s} \alpha_{sk}^{(i)} - \sum_{k=n_s+1}^{p_s} \alpha_{sk}^{(i)} \right) - \left( \sum_{k=1}^{q_2} \beta_{2k}^{(i)} + \sum_{k=1}^{q_3} \beta_{3k}^{(i)} + \dots + \sum_{k=1}^{q_s} \beta_{sk}^{(i)} \right) - \mu_i - \rho_i \\
&- \sum_{j=1}^k \lambda_j^{(i)} > 0 \quad (i = 1, \dots, r)
\end{aligned}$$

$$\text{(F)} \quad \left| \arg \left( z_i \prod_{j=1}^k (f_j t + g_j)^{\lambda_j^{(i)}} \right) \right| < \frac{1}{2} \Omega_i \pi \quad (a \leq t \leq b; i = 1, \dots, r)$$

**(G)**  $P \leq Q + 1$ . The equality holds, when , in addition,

$$\text{either } P > Q \text{ and } \left| z_i \left( \prod_{j=1}^k (f_j t + g_j)^{\lambda_j^{(i)}} \right) \right|^{\frac{1}{Q-P}} < 1 \quad (a \leq t \leq b)$$

$$\text{or } P \leq Q \text{ and } \max_{1 \leq i \leq k} \left[ \left| \left( z_i \prod_{j=1}^k (f_j t + g_j)^{\lambda_j^{(i)}} \right) \right| \right] < 1 \quad (a \leq t \leq b)$$

### Remark

By the following similar procedure, the results of this document can be extended to product of any finite number of multivariable I-functions defined by Prasad [1].

## 5. Conclusion

In this paper we have evaluated a generalized Eulerian integral involving the product of two multivariable I-functions defined by Prasad [1] with general arguments. The formulae established in this paper is very general nature. Thus, the results established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

## REFERENCES

- [1] Y.N. Prasad , Multivariable I-function , Vijnana Parishad Anusandhan Patrika 29 ( 1986 ) , page 231-237.
- [2] Saigo M. and Saxena R.K. Unified fractional integral formulas for the multivariable H-function I. J.Fractional Calculus 15 (1999), page 91-107.
- [3] Srivastava H.M. And Garg M. Some integral involving a general class of polynomials and multivariable H-function. Rev. Roumaine Phys. 32(1987), page 685-692.
- [4] Srivastava H.M. and Karlsson P.W. Multiple Gaussian Hypergeometric series. Ellis.Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.

[5] Srivastava H.M. and Manocha H.L : A treatise of generating functions. Ellis. Horwood.Series. Mathematics and Applications 1984, page 60

[6] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.