

Selberg integral involving a general sequence of functions, a class of polynomials and multivariable Aleph-functions

F.Y. AYANT¹

¹ Teacher in High School , France
 E-mail : fredericayant@gmail.com

ABSTRACT

In the present paper we evaluate the Selberg integral involving the product of a general sequence of functions, multivariable Aleph-functions and general class of polynomials of several variables. The importance of the result established in this paper lies in the fact they involve the Aleph-function of several variables which is sufficiently general in nature and capable to yielding a large of results merely by specializing the parameters their in.

Keywords:Multivariable Aleph-function, general class of polynomials, Selberg integral, General sequence of functions

2010 Mathematics Subject Classification. 33C99, 33C60, 44A20

1.Introduction and preliminaries.

The function Aleph of several variables generalize the multivariable I-function recently study by C.K. Sharma and Ahmad [6] , itself is an a generalisation of G and H-functions of multiple variables. The multiple Mellin-Barnes integral occuring in this paper will be referred to as the multivariables Aleph-function throughout our present study and will be defined and represented as follows.

$$\text{We define : } \mathfrak{N}(z_1, \dots, z_r) = \mathfrak{N}^{0, n; m_1, n_1, \dots, m_r, n_r}_{p_i, q_i, \tau_i; R; p_i(1), q_i(1), \tau_i(1); R^{(1)}; \dots; p_i(r), q_i(r), \tau_i(r); R^{(r)}} \left(\begin{array}{c} y_1 \\ \cdot \\ \cdot \\ \cdot \\ y_r \end{array} \right)$$

$$[(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, n}] \quad , [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1, p_i}] :$$

$$\dots \dots \dots \quad , [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1, q_i}] :$$

$$\left[(c_j^{(1)}; \gamma_j^{(1)})_{1, n_1}, [\tau_{i(1)}(c_{ji(1)}^{(1)}; \gamma_{ji(1)}^{(1)})_{n_1+1, p_i(1)}]; \dots ; [(c_j^{(r)}; \gamma_j^{(r)})_{1, n_r}], [\tau_{i(r)}(c_{ji(r)}^{(r)}; \gamma_{ji(r)}^{(r)})_{n_r+1, p_i(r)}] \right]$$

$$\left[(d_j^{(1)}; \delta_j^{(1)})_{1, m_1}, [\tau_{i(1)}(d_{ji(1)}^{(1)}; \delta_{ji(1)}^{(1)})_{m_1+1, q_i(1)}]; \dots ; [(d_j^{(r)}; \delta_j^{(r)})_{1, m_r}], [\tau_{i(r)}(d_{ji(r)}^{(r)}; \delta_{ji(r)}^{(r)})_{m_r+1, q_i(r)}] \right]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) y_k^{s_k} ds_1 \dots ds_r \tag{1.1}$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^n \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} s_k)]} \tag{1.2}$$

$$\text{and } \theta_k(s_k) = \frac{\prod_{j=1}^{m_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{n_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i(k)=1}^{R^{(k)}} [\tau_{i(k)} \prod_{j=m_k+1}^{q_i(k)} \Gamma(1 - d_{ji(k)}^{(k)} + \delta_{ji(k)}^{(k)} s_k) \prod_{j=n_k+1}^{p_i(k)} \Gamma(c_{ji(k)}^{(k)} - \gamma_{ji(k)}^{(k)} s_k)]} \tag{1.3}$$

Suppose , as usual , that the parameters

$$a_j, j = 1, \dots, p; b_j, j = 1, \dots, q;$$

$$c_j^{(k)}, j = 1, \dots, n_k; c_{j i^{(k)}}^{(k)}, j = n_k + 1, \dots, p_{i^{(k)}};$$

$$d_j^{(k)}, j = 1, \dots, m_k; d_{j i^{(k)}}^{(k)}, j = m_k + 1, \dots, q_{i^{(k)}};$$

$$\text{with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$$

are complex numbers , and the α' s, β' s, γ' s and δ' s are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{j i}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{j i^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{j i}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{j i^{(k)}}^{(k)} \leq 0 \quad (1.4)$$

The reals numbers τ_i are positives for $i = 1$ to R , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop , if necessary ,ensure that the poles of $\Gamma(d_j^{(k)} - \delta_j^{(k)} s_k)$ with $j = 1$ to m_k are separated from those of $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$ with $j = 1$ to n and $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$ with $j = 1$ to n_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{j i}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{j i}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{j i^{(k)}}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{j i^{(k)}}^{(k)} > 0, \text{ with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \quad (1.5)$$

The complex numbers z_i are not zero. Throughout this document , we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the the asymptotic expansion in the following convenient form :

$$\aleph(y_1, \dots, y_r) = O(|y_1|^{\alpha_1} \dots |y_r|^{\alpha_r}), \max(|y_1| \dots |y_r|) \rightarrow 0$$

$$\aleph(y_1, \dots, y_r) = O(|y_1|^{\beta_1} \dots |y_r|^{\beta_r}), \min(|y_1| \dots |y_r|) \rightarrow \infty$$

where, with $k = 1, \dots, r : \alpha_k = \min[Re(d_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, m_k$ and

$$\beta_k = \max[Re((c_j^{(k)} - 1) / \gamma_j^{(k)})], j = 1, \dots, n_k$$

Series representation of Aleph-function of several variables is given by

$$\aleph(y_1, \dots, y_r) = \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \dots \sum_{g_r=0}^{m_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1}^{G_1!} \dots \delta_{g_r}^{G_r!}} \psi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r})$$

$$\times \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) y_1^{-\eta_{G_1, g_1}} \dots y_r^{-\eta_{G_r, g_r}} \quad (1.6)$$

Where $\psi(\dots), \theta_i(\dots), i = 1, \dots, r$ are given respectively in (1.2), (1.3) and

$$\eta_{G_1, g_1} = \frac{d_{g_1}^{(1)} + G_1}{\delta_{g_1}^{(1)}}, \dots, \eta_{G_r, g_r} = \frac{d_{g_r}^{(r)} + G_r}{\delta_{g_r}^{(r)}}$$

which is valid under the conditions $\delta_{g_i}^{(i)}[d_j^i + p_i] \neq \delta_j^{(i)}[d_{g_i}^i + G_i]$ (1.7)

for $j \neq m_i, m_i = 1, \dots, \eta_{G_i, g_i}; p_i, n_i = 0, 1, 2, \dots, ; y_i \neq 0, i = 1, \dots, r$ (1.8)

Consider the Aleph-function of s variables

$$\aleph(z_1, \dots, z_s) = \aleph_{P_i, Q_i, t_i; r': P_{i(1)}, Q_{i(1)}, t_{i(1)}; r^{(1)}; \dots; P_{i(s)}, Q_{i(s)}, t_{i(s)}; r^{(s)}} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_s \end{matrix} \right)$$

$$[(u_j; \mu_j^{(1)}, \dots, \mu_j^{(r')})_{1, N}] \quad , [t_i(u_{ji}; \mu_{ji}^{(1)}, \dots, \mu_{ji}^{(r')})_{N+1, P_i}] :$$

$$\dots \dots \dots \quad , [t_i(v_{ji}; v_{ji}^{(1)}, \dots, v_{ji}^{(r')})_{M+1, Q_i}] :$$

$$\left[(a_j^{(1)}; \alpha_j^{(1)})_{1, N_1}, [t_{i(1)}(a_{ji(1)}^{(1)}; \alpha_{ji(1)}^{(1)})_{N_1+1, P_i^{(1)}}]; \dots; [(a_j^{(s)}; \alpha_j^{(s)})_{1, N_s}], [t_{i(s)}(a_{ji(s)}^{(s)}; \alpha_{ji(s)}^{(s)})_{N_s+1, P_i^{(s)}}] \right]$$

$$\left[(b_j^{(1)}; \beta_j^{(1)})_{1, M_1}, [t_{i(1)}(b_{ji(1)}^{(1)}; \beta_{ji(1)}^{(1)})_{M_1+1, Q_i^{(1)}}]; \dots; [(b_j^{(s)}; \beta_j^{(s)})_{1, M_s}], [t_{i(s)}(b_{ji(s)}^{(s)}; \beta_{ji(s)}^{(s)})_{M_s+1, Q_i^{(s)}}] \right]$$

$$= \frac{1}{(2\pi\omega)^s} \int_{L_1} \dots \int_{L_r} \zeta(t_1, \dots, t_s) \prod_{k=1}^s \phi_k(t_k) z_k^{t_k} dt_1 \dots dt_s \quad (1.9)$$

with $\omega = \sqrt{-1}$

$$\zeta(t_1, \dots, t_s) = \frac{\prod_{j=1}^N \Gamma(1 - u_j + \sum_{k=1}^s \mu_j^{(k)} t_k)}{\sum_{i=1}^{r'} [t_i \prod_{j=N+1}^{P_i} \Gamma(u_{ji} - \sum_{k=1}^s \mu_{ji}^{(k)} t_k) \prod_{j=1}^{Q_i} \Gamma(1 - v_{ji} + \sum_{k=1}^s v_{ji}^{(k)} t_k)]} \quad (1.10)$$

and $\phi_k(t_k) = \frac{\prod_{j=1}^{M_k} \Gamma(b_j^{(k)} - \beta_j^{(k)} t_k) \prod_{j=1}^{N_k} \Gamma(1 - a_j^{(k)} + \alpha_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{r^{(k)}} [t_i^{(k)} \prod_{j=M_k+1}^{Q_i^{(k)}} \Gamma(1 - b_{ji^{(k)}}^{(k)} + \beta_{ji^{(k)}}^{(k)} t_k) \prod_{j=N_k+1}^{P_i^{(k)}} \Gamma(a_{ji^{(k)}}^{(k)} - \alpha_{ji^{(k)}}^{(k)} s_k)]} \quad (1.11)$

Suppose, as usual, that the parameters

$u_j, j = 1, \dots, P; v_j, j = 1, \dots, Q;$

$a_j^{(k)}, j = 1, \dots, N_k; a_{j i^{(k)}}^{(k)}, j = n_k + 1, \dots, P_{i^{(k)}};$

$b_{j i^{(k)}}^{(k)}, j = m_k + 1, \dots, Q_{i^{(k)}}; b_j^{(k)}, j = 1, \dots, M_k;$

with $k = 1 \dots, s, i = 1, \dots, r', i^{(k)} = 1, \dots, r^{(k)}$

are complex numbers, and the $\alpha's, \beta's, \gamma's$ and $\delta's$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^N \mu_j^{(k)} + \iota_i \sum_{j=N+1}^{P_i} \mu_{ji}^{(k)} + \sum_{j=1}^{N_k} \alpha_j^{(k)} + \iota_{i^{(k)}} \sum_{j=N_k+1}^{P_{i^{(k)}}} \alpha_{j i^{(k)}}^{(k)} - \iota_i \sum_{j=1}^{Q_i} \nu_{ji}^{(k)} - \sum_{j=1}^{M_k} \beta_j^{(k)} - \iota_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \beta_{j i^{(k)}}^{(k)} \leq 0 \quad (1.12)$$

The reals numbers τ_i are positives for $i = 1, \dots, r$, $\iota_{i^{(k)}}$ are positives for $i^{(k)} = 1 \dots r^{(k)}$

The contour L_k is in the t_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary, ensure that the poles of $\Gamma(b_j^{(k)} - \beta_j^{(k)} t_k)$ with $j = 1$ to M_k are separated from those of $\Gamma(1 - u_j + \sum_{i=1}^s \mu_j^{(k)} t_k)$ with $j = 1$ to N and $\Gamma(1 - a_j^{(k)} + \alpha_j^{(k)} t_k)$ with $j = 1$ to N_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} B_i^{(k)} \pi, \text{ where}$$

$$B_i^{(k)} = \sum_{j=1}^N \mu_j^{(k)} - \iota_i \sum_{j=N+1}^{P_i} \mu_{ji}^{(k)} - \iota_i \sum_{j=1}^{Q_i} \nu_{ji}^{(k)} + \sum_{j=1}^{N_k} \alpha_j^{(k)} - \iota_{i^{(k)}} \sum_{j=N_k+1}^{P_{i^{(k)}}} \alpha_{j i^{(k)}}^{(k)} + \sum_{j=1}^{M_k} \beta_j^{(k)} - \iota_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \beta_{j i^{(k)}}^{(k)} > 0, \text{ with } k = 1 \dots, s, i = 1, \dots, r, i^{(k)} = 1, \dots, r^{(k)} \quad (1.13)$$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_s) = O(|z_1|^{\alpha'_1} \dots |z_s|^{\alpha'_s}), \max(|z_1| \dots |z_s|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_s) = O(|z_1|^{\beta'_1} \dots |z_s|^{\beta'_s}), \min(|z_1| \dots |z_s|) \rightarrow \infty$$

where, with $k = 1, \dots, s, z : \alpha'_k = \min[Re(b_j^{(k)} / \beta_j^{(k)})], j = 1, \dots, M_k$ and

$$\beta'_k = \max[Re((a_j^{(k)} - 1) / \alpha_j^{(k)})], j = 1, \dots, N_k$$

We will use these following notations in this paper

$$U = P_i, Q_i, \ell_i; r'; V = M_1, N_1; \dots; M_s, N_s \quad (1.15)$$

$$W = P_{i(1)}, Q_{i(1)}, \ell_{i(1)}; r^{(1)}, \dots, P_{i(r)}, Q_{i(r)}, \ell_{i(s)}; r^{(s)} \quad (1.16)$$

$$A = \{(u_j; \mu_j^{(1)}, \dots, \mu_j^{(s)})_{1,N}\}, \{\ell_i(u_{ji}; \mu_{ji}^{(1)}, \dots, \mu_{ji}^{(s)})_{N+1, P_i}\} \quad (1.17)$$

$$B = \{\ell_i(v_{ji}; v_{ji}^{(1)}, \dots, v_{ji}^{(s)})_{M+1, Q_i}\} \quad (1.18)$$

$$C = (a_j^{(1)}; \alpha_j^{(1)})_{1, N_1}, \ell_{i(1)}(a_{ji(1)}^{(1)}; \alpha_{ji(1)}^{(1)})_{N_1+1, P_{i(1)}}, \dots, (a_j^{(s)}; \alpha_j^{(s)})_{1, N_s}, \ell_{i(s)}(a_{ji(s)}^{(s)}; \alpha_{ji(s)}^{(s)})_{N_s+1, P_{i(s)}} \quad (1.19)$$

$$D = (b_j^{(1)}; \beta_j^{(1)})_{1, M_1}, \ell_{i(1)}(b_{ji(1)}^{(1)}; \beta_{ji(1)}^{(1)})_{M_1+1, Q_{i(1)}}, \dots, (b_j^{(s)}; \beta_j^{(s)})_{1, M_s}, \ell_{i(s)}(\beta_{ji(s)}^{(s)}; \beta_{ji(s)}^{(s)})_{M_s+1, Q_{i(s)}} \quad (1.20)$$

The multivariable Aleph-function write :

$$\aleph(z_1, \dots, z_s) = \aleph_{U:W}^{0, N:V} \left(\begin{array}{c|c} z_1 & \text{A : C} \\ \cdot & \cdot \cdot \cdot \\ \cdot & \text{B : D} \\ z_s & \end{array} \right) \quad (1.21)$$

The generalized polynomials defined by Srivastava [9], is given in the following manner :

$$S_{N_1, \dots, N_t}^{M_1, \dots, M_t} [y_1, \dots, y_t] = \sum_{K_1=0}^{[N_1/M_1]} \dots \sum_{K_t=0}^{[N_t/M_t]} \frac{(-N_1)_{M_1 K_1}}{K_1!} \dots \frac{(-N_t)_{M_t K_t}}{K_t!}$$

$$A[N_1, K_1; \dots; N_t, K_t] y_1^{K_1} \dots y_t^{K_t} \quad (1.22)$$

Where M_1, \dots, M_s are arbitrary positive integers and the coefficients $A[N_1, K_1; \dots; N_t, K_t]$ are arbitrary constants, real or complex. In the present paper, we use the following notation

$$a_1 = \frac{(-N_1)_{M_1 K_1}}{K_1!} \dots \frac{(-N_t)_{M_t K_t}}{K_t!} A[N_1, K_1; \dots; N_t, K_t] \quad (1.23)$$

In the document , we note :

$$G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) = \phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) \quad (1.24)$$

where $\phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}), \theta_1(\eta_{G_1, g_1}), \dots, \theta_r(\eta_{G_r, g_r})$ are given respectively in (1.2) and (1.3)

2. Sequence of function

Agarwal and Chaubey [1], Salim [5] and several others have studied a general sequence of functions. In the present document we shall study the following useful series formula for a general sequence of functions.

$$R_n^{\alpha, \beta} [x; E, F, g, h; p, q; \gamma; \delta; e^{sx^r}] = \sum_{w, v, u, t, e, k_1, k_2} \psi(w, v, u, t, e, k_1, k_2) x^R \quad (2.1)$$

$$\text{where } \sum_{w, v, u, t, e, k_1, k_2} = \sum_{w=0}^{\infty} \sum_{v=0}^n \sum_{u=0}^v \sum_{t=0}^n \sum_{c=0}^t \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \quad (2.2)$$

and the infinite series on the right side (2.1) is absolutely convergent, $R = ln + qv + pt + rw + k_1r + k_2q$

$$\text{and } \psi(w, v, u, t, e, k_1, k_2) = \frac{(-)^{t+w+k_2}(-v)_u(-t)_e(\alpha)_t l^n}{w!v!u!t!e!K_n k_1!k_2!} \frac{s^{w+k_1} F^{\gamma n-t}}{(1-\alpha-t)_e} (\alpha-\gamma n)_e$$

$$(-\beta-\delta n)_v g^{v+k_2} h^{\delta n-v-k_2} (v-\delta n)_{k_2} E^t \left(\frac{pe+rw+\lambda+qn}{l} \right)_n \tag{2.3}$$

where K_n is a sequence of constants.

By suitably specializing the parameters involving in (2.1), a general sequence of function reduced to generalized polynomials set studied by Raizada [4], a class of polynomials introduced by Fujiwara [3] and several others authors.

3. Required integral

We have the following result, see (Beals et al [2], page 54)

$$S(a, b, c) = \int_0^1 \cdots \int_0^1 \prod_{i=1}^n x_i^{a-1} (1-x_i)^{b-1} \prod_{1 \leq j < k \leq n} |x_j - x_k|^{2c} dx_1 \cdots dx_n =$$

$$= \prod_{j=0}^{n-1} \frac{\Gamma(a+jc)\Gamma(b+jc)\Gamma(1+(j+1)c)}{\Gamma(a+b+(n-1+j)c)\Gamma(1+c)} \tag{3.1}$$

with $Re(a) > 0, Re(b) > 0, Re(c) > Max \left\{ -\frac{1}{n}, -\frac{Re(a)}{n-1}, -\frac{Re(b)}{n-1} \right\}$

$S(a, b, c)$ is the Selberg integral with three parameters

4. Main integral

Let $X_{u,v,w} = \prod_{i=1}^n x_i^u (1-x_i)^v \prod_{1 \leq j < k \leq n} |x_j - x_k|^{2w}$

we have the following formula

Theorem

$$\int_0^1 \cdots \int_0^1 \prod_{i=1}^n x_i^{a-1} (1-x_i)^{b-1} \prod_{1 \leq j < k \leq n} |x_j - x_k|^{2c} R_n^{\alpha,\beta} [z X_{\alpha',\beta',\gamma'}; E, F, g, h; p, q; \gamma; \delta; e^{s(z X_{\alpha',\beta',\gamma'})^r}]$$

$$S_{N_1, \dots, N_t}^{M_1, \dots, M_t} \left(\begin{matrix} y_1 X_{\alpha_1, \beta_1, \gamma_1} \\ \dots \\ y_t X_{\alpha_t, \beta_t, \gamma_t} \end{matrix} \right) \mathfrak{N}_{u:w}^{0,n:v} \left(\begin{matrix} z_1 X_{\delta_1, \psi_1, \phi_1} \\ \dots \\ z_r X_{\delta_r, \psi_r, \phi_r} \end{matrix} \right) \mathfrak{N}_{U:W}^{0,N:V} \left(\begin{matrix} Z_1 X_{\epsilon_1, \eta_1, \zeta_1} \\ \dots \\ Z_s X_{\epsilon_s, \eta_s, \zeta_s} \end{matrix} \right) dx_1 \cdots dx_n =$$

$$\sum_{w,v,u,t,e,k_1,k_2} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \cdots \sum_{g_r=0}^{m_r} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_t=0}^{[N_t/M_t]} a_1 \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \cdots \delta_{g_r} G_r!} \psi(w, v, u, t, e, k_1, k_2)$$

$$\begin{aligned}
& G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) y_1^{K_1} \dots y_t^{K_t} z_1^{\eta_{G_1, g_1}} \dots z_r^{\eta_{G_r, g_r}} z^R \mathbb{N}_{U_{3n, 1+n}: W}^{0, N+3n: V} \left(\begin{array}{c} Z_1 \\ \dots \\ Z_s \end{array} \right) \\
& [1-a-\alpha'R - \sum_{i=1}^t K_i \alpha_i - \sum_{i=1}^r \eta_{G_i, g_i} \delta_i - j(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), \epsilon_1 + j\zeta_1, \dots, \epsilon_s + j\zeta_s]_{0, n-1} \\
& \quad \quad \quad \dots \\
& \quad \quad \quad (-c-\gamma'R + \sum_{i=1}^t K_i \gamma_i - \sum_{i=1}^r \phi_i \eta_{G_i, g_i}, \zeta_1, \dots, \zeta_s), \\
& [1-b-\beta'R - \sum_{i=1}^t K_i \beta_i - \sum_{i=1}^r \eta_{G_i, g_i} \psi_i - j(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), \eta_1 + j\zeta_1, \dots, \eta_s + j\zeta_s]_{0, n-1} \\
& \quad \quad \quad \dots \\
& \quad \quad \quad B_1 \\
& -(j+1)(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), j\zeta_1, \dots, j\zeta_s]_{0, n-1}, A : C \\
& \quad \quad \quad \dots \\
& \quad \quad \quad B : D
\end{aligned} \tag{4.1}$$

where $B_1 = [1 - a - b - (\alpha' + \beta')R - \sum_{i=1}^t K_i(\alpha_i + \beta_i) - \sum_{i=1}^r (\delta_i + \phi_i)\eta_{G_i, g_i} - (n - 1 + j) \times (c + \gamma'R + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}); \epsilon_1 + \eta_1 + j\zeta_1, \dots, \epsilon_s + \eta_s + j\zeta_s]_{0, n-1}$ and $U_{3n, n+1} = P_i + 3n; Q_i + n + 1; \nu_i; r'$

Provided that

a) $\min\{\alpha', \beta', \gamma', \alpha_i, \beta_i, \gamma_i, \delta_j, \psi_j, \phi_j, \epsilon_k, \eta_k, \zeta_k\} > 0, i = 1, \dots, t, j = 1, \dots, r, k = 1, \dots, s,$

b) $A = Re[a + \alpha'R + \sum_{i=1}^r \delta_i \min_{1 \leq j \leq m_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^s \epsilon_i \min_{1 \leq j \leq M_i} \frac{b_j^{(i)}}{\beta_j^{(i)}}] > 0$

c) $B = Re[b + \beta'R + \sum_{i=1}^r \psi_i \min_{1 \leq j \leq m_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^s \eta_i \min_{1 \leq j \leq M_i} \frac{b_j^{(i)}}{\beta_j^{(i)}}] > 0$

d) $C = Re[c + \gamma'R + \sum_{i=1}^r \phi_i \min_{1 \leq j \leq m_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^s \zeta_i \min_{1 \leq j \leq M_i} \frac{b_j^{(i)}}{\beta_j^{(i)}}] > Max \left\{ -\frac{1}{n}, -\frac{A}{n-1}, -\frac{B}{n-1} \right\}$

e) $|arg z_k| < \frac{1}{2} A_i^{(k)} \pi,$ where $A_i^{(k)}$ is defined by (1.5); $i = 1, \dots, r$

f) $|arg Z_k| < \frac{1}{2} B_i^{(k)} \pi,$ where $B_i^{(k)}$ is defined by (1.13); $i = 1, \dots, s$

Proof

under the same notations and conditions that (4.1)

6. Aleph-function of two variables

If $s = 2$, we obtain the Aleph-function of two variables defined by K.Sharma [8], and we have the following simple integrals.

Corollary 2

$$\int_0^1 \cdots \int_0^1 \prod_{i=1}^n x_i^{a-1} (1-x_i)^{b-1} \prod_{1 \leq j < k \leq n} |x_j - x_k|^{2c} R_n^{\alpha, \beta} [z X_{\alpha', \beta', \gamma'}; E, F, g, h; p, q; \gamma; \delta; e^{s(z X_{\alpha', \beta', \gamma'})^r}]$$

$$S_{N_1, \dots, N_t}^{M_1, \dots, M_t} \left(\begin{matrix} y_1 X_{\alpha_1, \beta_1, \gamma_1} \\ \dots \\ y_t X_{\alpha_t, \beta_t, \gamma_t} \end{matrix} \right) \aleph_{u:w}^{0, n:v} \left(\begin{matrix} z_1 X_{\delta_1, \psi_1, \phi_1} \\ \dots \\ z_r X_{\delta_r, \psi_r, \phi_r} \end{matrix} \right) \aleph_{U:W}^{0, N:V} \left(\begin{matrix} Z_1 X_{\epsilon_1, \eta_1, \zeta_1} \\ \dots \\ Z_2 X_{\epsilon_2, \eta_2, \zeta_2} \end{matrix} \right) dx_1 \cdots dx_n =$$

$$\sum_{w, v, u, t, e, k_1, k_2} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \cdots \sum_{g_r=0}^{m_r} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_t=0}^{[N_t/M_t]} a_1 \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \cdots \delta_{g_r} G_r!} \psi(w, v, u, t, e, k_1, k_2)$$

$$G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) y_1^{K_1} \cdots y_t^{K_t} z_1^{\eta_{G_1, g_1}} \cdots z_r^{\eta_{G_r, g_r}} z^R \aleph_{U_{3n, 1+n}:W}^{0, N+3n:V} \left(\begin{matrix} Z_1 \\ \dots \\ Z_2 \end{matrix} \right)$$

$$[1-a-\alpha'R - \sum_{i=1}^t K_i \alpha_i - \sum_{i=1}^r \eta_{G_i, g_i} \delta_i - j(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), \epsilon_1 + j\zeta_1, \epsilon_2 + j\zeta_2]_{0, n-1}$$

$$\dots$$

$$(-c-\gamma'R + \sum_{i=1}^t K_i \gamma_i - \sum_{i=1}^r \phi_i \eta_{G_i, g_i}, \zeta_1, \zeta_2),$$

$$[1-b-\beta'R - \sum_{i=1}^t K_i \beta_i - \sum_{i=1}^r \eta_{G_i, g_i} \psi_i - j(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), \eta_1 + j\zeta_1, \eta_2 + j\zeta_2]_{0, n-1}$$

$$\dots$$

$$B_1$$

$$-(j+1)(c+\gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), j\zeta_1, j\zeta_2]_{0, n-1}, A : C$$

$$\dots$$

$$B : D \quad (6.1)$$

where $B_1 = [1 - a - b - (\alpha' + \beta')R - \sum_{i=1}^t K_i(\alpha_i + \beta_i) - \sum_{i=1}^r (\delta_i + \phi_i)\eta_{G_i, g_i} - (n - 1 + j)c \times$

$$(c + \gamma'R + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}); \epsilon_1 + \eta_1 + j\zeta_1, \epsilon_2 + \eta_2 + j\zeta_2]_{0, n-1}$$

and $U_{3n, n+1} = P_i + 3n; Q_i + n + 1; \iota_i; r'$

under the same notations and conditions that (4.1) with $s = 2$

7. I-function of two variables

If $\iota_i, \iota'_i, \iota''_i \rightarrow 1$, then the Aleph-function of two variables degenerates in the I-function of two variables defined by sharma et al [7] and we obtain the same formula with the I-function of two variables.

Corollary 3

$$\int_0^1 \cdots \int_0^1 \prod_{i=1}^n x_i^{a-1} (1-x_i)^{b-1} \prod_{1 \leq j < k \leq n} |x_j - x_k|^{2c} R_n^{\alpha, \beta} [z X_{\alpha', \beta', \gamma'}; E, F, g, h; p, q; \gamma; \delta; e^{s(z X_{\alpha', \beta', \gamma'})^r}]$$

$$S_{N_1, \dots, N_t}^{M_1, \dots, M_t} \left(\begin{matrix} y_1 X_{\alpha_1, \beta_1, \gamma_1} \\ \dots \\ y_t X_{\alpha_t, \beta_t, \gamma_t} \end{matrix} \right) X_{u:w}^{0, n:v} \left(\begin{matrix} z_1 X_{\delta_1, \psi_1, \phi_1} \\ \dots \\ z_r X_{\delta_r, \psi_r, \phi_r} \end{matrix} \right) I_{U:W}^{0, N:V} \left(\begin{matrix} Z_1 X_{\epsilon_1, \eta_1, \zeta_1} \\ \dots \\ Z_s X_{\epsilon_2, \eta_2, \zeta_2} \end{matrix} \right) dx_1 \cdots dx_n =$$

$$\sum_{w, v, u, t, e, k_1, k_2} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \cdots \sum_{g_r=0}^{m_r} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_t=0}^{[N_t/M_t]} a_1 \frac{(-)^{G_1 + \dots + G_r}}{\delta_{g_1} G_1! \cdots \delta_{g_r} G_r!} \psi(w, v, u, t, e, k_1, k_2)$$

$$G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) y_1^{K_1} \cdots y_t^{K_t} z_1^{\eta_{G_1, g_1}} \cdots z_r^{\eta_{G_r, g_r}} z^R I_{U_{3n, 1+n}:W}^{0, N+3n:V} \left(\begin{matrix} Z_1 \\ \dots \\ Z_2 \end{matrix} \right)$$

$$[1-a-\alpha'R - \sum_{i=1}^t K_i \alpha_i - \sum_{i=1}^r \eta_{G_i, g_i} \delta_i - j(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), \epsilon_1 + j\zeta_1, \epsilon_2 + j\zeta_2]_{0, n-1}$$

$$\dots$$

$$(-c-\gamma'R + \sum_{i=1}^t K_i \gamma_i - \sum_{i=1}^r \phi_i \eta_{G_i, g_i}, \zeta_1, \zeta_2),$$

$$[1-b-\beta'R - \sum_{i=1}^t K_i \beta_i - \sum_{i=1}^r \eta_{G_i, g_i} \psi_i - j(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}), \eta_1 + j\zeta_1, \eta_2 + j\zeta_2]_{0, n-1}$$

$$\dots$$

$$B_1$$

$$-(j+1)(c + \gamma'R + \sum_{i=1}^t \gamma_i K_i + \sum_{i=1}^r \phi_i \eta_{G_i, g_i}, j\zeta_1, j\zeta_2)_{0, n-1}, A : C$$

$$\dots$$

$$B : D \tag{7.1}$$

under the same conditions and notation that (4.1) with $s = 2$ and $\iota_i, \iota'_i, \iota''_i \rightarrow 1$

8. Conclusion

In this paper we have evaluated a Selberg integral involving the multivariable Aleph-functions, a class of polynomials of several variables and the general of sequence of functions. The integral established in this paper is of very general nature as it contains Multivariable Aleph-function, which is a general function of several variables studied so far. Thus, the integral established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

- [1] Agrawal B.D. And Chaubey J.P. Certain derivation of generating relations for generalized polynomials. Indian J. Pure and Appl. Math 10 (1980), page 1155-1157, ibid 11 (1981), page 357-359
- [2] Beals Richard and Roderick Wong. Special functions. Cambridge University Press 2010
- [3] Fujiwara I. A unified presentation of classical orthogonal polynomials. Math. Japon. 11 (1966), page133-148.
- [4] Raizada S.K. A study of unified representation of special functions of Mathematics Physics and their use in statistical and boundary value problem. Ph.D. Thesis, Bundelkhand University, Jhansi, India, 1991
- [5] Salim T.O. A serie formula of generalized class of polynomials associated with Laplace transform and fractional integral operators. J. Rajasthan. Acad. Phy. Sci. 1(3) (2002), page 167-176.
- [6] Sharma C.K.and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20,no2, p 113-116.
- [7] C.K. Sharma and P.L. mishra : On the I-function of two variables and its properties. Acta Ciencia Indica Math , 1991, Vol 17 page 667-672.
- [8] Sharma K. On the integral representation and applications of the generalized function of two variables , International Journal of Mathematical Engineering and Sciences , Vol 3 , issue1 (2014) , page1-13.
- [9] Srivastava H.M. A multilinear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial, Pacific. J. Math. 177(1985), page183-191.