

# On general multiple Eulerian integrals involving the multivariable Aleph-function , a general class of polynomials and the Aleph-function of one variable

F.Y. AYANT<sup>1</sup>

<sup>1</sup> Teacher in High School , France  
 E-mail : fredericayant@gmail.com

## ABSTRACT

The object of this paper is first, to evaluate a general multiple Eulerian integrals with general integrands involving the product of a multivariable Aleph-function , a general class of polynomials, the Aleph-function of one variable and generalized hypergeometric function with general arguments. The second multiple Eulerian integral contain a general class of polynomials, the general polynomial set, the Aleph-function of one variable and multivariable Aleph-function with general arguments. Our integral formulas are interesting and unified nature.

Keywords :Multivariable Aleph-function, class of polynomial, general polynomials set, multiple Eulerian integral, Generalized incomplete hypergeometric function, multivariable I-function.

**2010 Mathematics Subject Classification.** 33C99, 33C60, 44A20

## 1. Introduction

In this paper, we evaluate two multiple Eulerian integrals involving the multivariable Aleph-function , the Aleph-function of one variable and multivariable class of polynomials with general arguments.

The Aleph-function of several variables generalize the multivariable I-function defined by Sharma and Ahmad [6] , itself is an a generalisation of G and H-functions of several variables. The multiple Mellin-Barnes integral occurring in this paper will be referred to as the multivariable Aleph-function throughout our present study and will be defined and represented as follows.

$$\text{We have : } \aleph(z_1, \dots, z_r) = \aleph_{p_i, q_i, \tau_i; R: p_i(1), q_i(1), \tau_i(1); R^{(1)}; \dots; p_i(r), q_i(r), \tau_i(r); R^{(r)}}^{0, n: m_1, n_1, \dots, m_r, n_r} \left( \begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \right)$$

$$[(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, n}] \quad , [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1, p_i}] :$$

$$\dots\dots\dots, [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1, q_i}] :$$

$$\left( \begin{matrix} [(c_j^{(1)}), \gamma_j^{(1)})_{1, n_1}], [\tau_{i(1)}(c_{ji(1)}^{(1)}, \gamma_{ji(1)}^{(1)})_{n_1+1, p_i^{(1)}}]; \dots ; [(c_j^{(r)}), \gamma_j^{(r)})_{1, n_r}], [\tau_{i(r)}(c_{ji(r)}^{(r)}, \gamma_{ji(r)}^{(r)})_{n_r+1, p_i^{(r)}}] \\ [(d_j^{(1)}), \delta_j^{(1)})_{1, m_1}], [\tau_{i(1)}(d_{ji(1)}^{(1)}, \delta_{ji(1)}^{(1)})_{m_1+1, q_i^{(1)}}]; \dots ; [(d_j^{(r)}), \delta_j^{(r)})_{1, m_r}], [\tau_{i(r)}(d_{ji(r)}^{(r)}, \delta_{ji(r)}^{(r)})_{m_r+1, q_i^{(r)}}] \end{matrix} \right)$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} ds_1 \dots ds_r \quad (1.1)$$

with  $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^n \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} s_k)]} \quad (1.2)$$

$$\text{and } \theta_k(s_k) = \frac{\prod_{j=1}^{m_k} \Gamma(d_j^{(k)} - \delta_j'^{(k)} s_k) \prod_{j=1}^{n_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=m_k+1}^{q_{i^{(k)}}} \Gamma(1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}'^{(k)} s_k) \prod_{j=n_k+1}^{p_{i^{(k)}}} \Gamma(c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} s_k)]} \quad (1.3)$$

Suppose , as usual , that the parameters

$$a_j, j = 1, \dots, p; b_j, j = 1, \dots, q;$$

$$c_j^{(k)}, j = 1, \dots, n_k; c_{ji^{(k)}}^{(k)}, j = n_k + 1, \dots, p_{i^{(k)}};$$

$$d_j^{(k)}, j = 1, \dots, m_k; d_{ji^{(k)}}^{(k)}, j = m_k + 1, \dots, q_{i^{(k)}};$$

$$\text{with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$$

are complex numbers , and the  $\alpha' s, \beta' s, \gamma' s$  and  $\delta' s$  are assumed to be positive real numbers for standardization purpose such that

$$\begin{aligned} U_i^{(k)} &= \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j'^{(k)} \\ &\quad - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}'^{(k)} \leq 0 \end{aligned} \quad (1.4)$$

The reals numbers  $\tau_i$  are positives for  $i = 1$  to  $R$  ,  $\tau_{i^{(k)}}$  are positives for  $i^{(k)} = 1$  to  $R^{(k)}$

The contour  $L_k$  is in the  $s_k$ -p lane and run from  $\sigma - i\infty$  to  $\sigma + i\infty$  where  $\sigma$  is a real number with loop , if necessary , ensure that the poles of  $\Gamma(d_j^{(k)} - \delta_j'^{(k)} s_k)$  with  $j = 1$  to  $m_k$  are separated from those of  $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$  with  $j = 1$  to  $n$  and  $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$  with  $j = 1$  to  $n_k$  to the left of the contour  $L_k$  . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$\begin{aligned} A_i^{(k)} &= \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} \\ &+ \sum_{j=1}^{m_k} \delta_j'^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}'^{(k)} > 0, \text{ with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \end{aligned} \quad (1.5)$$

The complex numbers  $z_i$  are not zero. Throughout this document , we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the the asymptotic expansion in the following convenient form :

$$\aleph(y_1, \dots, y_r) = O(|y_1|^{\alpha_1}, \dots, |y_r|^{\alpha_r}), \max(|y_1|, \dots, |y_r|) \rightarrow 0$$

$$\aleph(y_1, \dots, y_r) = O(|y_1|^{\beta_1}, \dots, |y_r|^{\beta_r}), \min(|y_1|, \dots, |y_r|) \rightarrow \infty$$

where  $k = 1, \dots, r : \alpha_k = \min[Re(d_j^{(k)} / \delta_j'^{(k)})], j = 1, \dots, m_k$  and

$$\beta_k = \max[Re((c_j^{(k)} - 1)/\gamma_j^{(k)})], j = 1, \dots, n_k$$

We will use these following notations in this paper

$$U = p_i, q_i, \tau_i; R; V = m_1, n_1; \dots; m_r, n_r \quad (1.6)$$

$$W = p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}, \dots, p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)} \quad (1.7)$$

$$A = \{(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1,n}\}, \{\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1, p_i}\} \quad (1.8)$$

$$B = \{\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1, q_i}\} \quad (1.9)$$

$$C = \{(c_j^{(1)}; \gamma_j^{(1)})_{1, n_1}\}, \tau_{i(1)}(c_{ji(1)}^{(1)}; \gamma_{ji(1)}^{(1)})_{n_1+1, p_{i(1)}}\}, \dots, \{(c_j^{(r)}; \gamma_j^{(r)})_{1, n_r}\}, \tau_{i(r)}(c_{ji(r)}^{(r)}; \gamma_{ji(r)}^{(r)})_{n_r+1, p_{i(r)}}\} \quad (1.10)$$

$$D = \{(d_j^{(1)}; \delta_j^{(1)})_{1, m_1}\}, \tau_{i(1)}(d_{ji(1)}^{(1)}; \delta_{ji(1)}^{(1)})_{m_1+1, q_{i(1)}}\}, \dots, \{(d_j^{(r)}; \delta_j^{(r)})_{1, m_r}\}, \tau_{i(r)}(d_{ji(r)}^{(r)}; \delta_{ji(r)}^{(r)})_{m_r+1, q_{i(r)}}\} \quad (1.11)$$

The contracted form is :

$$\aleph(z_1, \dots, z_r) = \aleph_{U:W}^{0, n; V} \left( \begin{array}{c|c} z_1 & A : C \\ \vdots & \\ \vdots & . \ . \ . \\ \vdots & B : D \\ z_r & \end{array} \right) \quad (1.12)$$

The Aleph- function , introduced by Südländ [10] et al , however the notation and complete definition is presented here in the following manner in terms of the Mellin-Barnes type integral :

$$\aleph(z) = \aleph_{P_i, Q_i, c_i; r'}^{M, N} \left( z \left| \begin{array}{l} (a_j, A_j)_{1, n}, [c_i(a_{ji}, A_{ji})]_{n+1, p_i; r} \\ (b_j, B_j)_{1, m}, [c_i(b_{ji}, B_{ji})]_{m+1, q_i; r} \end{array} \right. \right) = \frac{1}{2\pi\omega} \int_L \Omega_{P_i, Q_i, c_i; r}^{M, N}(s) z^{-s} ds \quad (1.13)$$

for all  $z$  different to 0 and

$$\Omega_{P_i, Q_i, c_i; r'}^{M, N}(s) = \frac{\prod_{j=1}^M \Gamma(b_j - B_j s) \prod_{j=1}^N \Gamma(1 - a_j + A_j s)}{\sum_{i=1}^r c_i \prod_{j=N+1}^{P_i} \Gamma(a_{ji} - A_{ji} s) \prod_{j=M+1}^{Q_i} \Gamma(1 - b_{ji} + B_{ji} s)} \quad (1.14)$$

$$\text{With : } |arg z| < \frac{1}{2}\pi\Omega \quad \text{Where } \Omega = \sum_{j=1}^M \beta_j + \sum_{j=1}^N \alpha_j - c_i \left( \sum_{j=M+1}^{Q_i} \beta_{ji} + \sum_{j=N+1}^{P_i} \alpha_{ji} \right) > 0; i = 1, \dots, r'$$

For convergence conditions and other details of Aleph-function , see Südländ et al [10], the serie representation of Aleph-function is given by Chaurasia et al [2].

$$\aleph_{P_i, Q_i, c_i; r'}^{M, N}(z) = \sum_{G=1}^M \sum_{g=0}^{\infty} \frac{(-)^g \Omega_{P_i, Q_i, c_i, r}^{M, N}(s)}{B_G g!} z^s \quad (1.15)$$

$$\text{With } s = \eta_{G, g} = \frac{b_G + g}{B_G}, P_i < Q_i, |z| < 1 \text{ and } \Omega_{P_i, Q_i, c_i; r}^{M, N}(s) \text{ is given in (1.2)} \quad (1.16)$$

The generalized polynomials defined by Srivastava [7], is given in the following manner :

$$S_{N_1, \dots, N_u}^{M_1, \dots, M_u} [y_1, \dots, y_u] = \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_u=0}^{[N_u/M_u]} \frac{(-N_1)_{M_1 K_1}}{K_1!} \cdots \frac{(-N_u)_{M_u K_u}}{K_u!} A[N_1, K_1; \dots; N_u, K_u] y_1^{K_1} \cdots y_u^{K_u} \quad (1.17)$$

Where  $M_1, \dots, M_u$  are arbitrary positive integers and the coefficients  $A[N_1, K_1; \dots; N_u, K_u]$  are arbitrary constants, real or complex. In the present paper, we use the following notation

$$a_u = \frac{(-N_1)_{M_1 K_1}}{K_1!} \cdots \frac{(-N_u)_{M_u K_u}}{K_u!} A[N_1, K_1; \dots; N_u, K_u] \quad (1.18)$$

The multivariable I-function is defined in term of multiple Mellin-Barnes type integral :

## 2. Sequence of functions

Agarwal and Chaubey [1], Salim [5] and several others have studied a general sequence of functions. In the present document we shall study the following useful series formula for a general sequence of functions.

$$R_n^{\alpha, \beta} [x; E, F, g, h; p, q; \gamma; \delta; e^{-sx^r}] = \sum_{w, v, u, t, e, k_1, k_2} \psi(w, v, u, t, e, k_1, k_2) x^R \quad (2.1)$$

$$\text{where } \sum_{w, v, u, t, e, k_1, k_2} = \sum_{w=0}^{\infty} \sum_{v=0}^n \sum_{u=0}^v \sum_{t=0}^n \sum_{c=0}^t \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \quad (2.2)$$

and the infinite series on the right side (2.1) is absolutely convergent,  $R = ln + qv + pt + rw + k_1 r + k_2 q$  (2.3)

$$\text{and } \psi(w, v, u, t, e, k_1, k_2) = \frac{(-)^{t+w+k_2} (-v)_u (-t)_e (\alpha)_t l^n}{w! v! u! t! e! K_n k_1! k_2!} \frac{s^{w+k_1} F^{\gamma n-t}}{(1-\alpha-t)_e} (\alpha - \gamma n)_e$$

$$(-\beta - \delta n)_v g^{v+k_2} h^{\delta n-v-k_2} (v - \delta n)_{k_2} E^t \left( \frac{pe + rw + \lambda + qn}{l} \right)_n \quad (2.4)$$

where  $K_n$  is a sequence of constants. This function will note  $R_n^{\alpha, \beta} [x]$

By suitably specializing the parameters involving in (2.1), a general sequence of function reduced to generalized polynomials set studied by Raizada [4], a class of polynomials introduced by Fujiwara [3] and several others authors.

## 3. Integral representation of generalized hypergeometric function

The following generalized hypergeometric function in terms of multiple contour integrals is also required [8 ,page 39 eq .30]

$$\begin{aligned} & \frac{\prod_{j=1}^P \Gamma(A_j)}{\prod_{j=1}^Q \Gamma(B_j)} {}_P F_Q [(A_P); (B_Q); -(x_1 + \cdots + x_r)] \\ &= \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \frac{\prod_{j=1}^P \Gamma(A_j + s_1 + \cdots + s_r)}{\prod_{j=1}^Q \Gamma(B_j + s_1 + \cdots + s_r)} \Gamma(-s_1) \cdots \Gamma(-s_r) x_1^{s_1} \cdots x_r^{s_r} ds_1 \cdots ds_r \end{aligned} \quad (3.1)$$

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of  $\Gamma(A_j + s_1 + \cdots + s_r)$  are separated from those of  $\Gamma(-s_j), j = 1, \dots, r$ . The above result (1.23) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of  $\Gamma(-s_j), j = 1, \dots, r$

The equivalent form of Eulerian beta integral is :

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} dt = (b-a)^{\alpha+\beta} B(\alpha, \beta) (Re(\alpha) > 0, Re(\beta) > 0, a \neq b) \quad (3.2)$$

#### 4. First integral

We note :

$$V_1 = V; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0; W_1 = W; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1 \quad (4.1)$$

$$C_1 = C; (1, 0), \dots, (1, 0); (1, 0), \dots, (1, 0); D_1 = D; (0, 1), \dots, (0, 1); (0, 1), \dots, (0, 1) \quad (4.2)$$

$$A^* = [1 + \sigma'_i - \eta_{G,g} c'_i - \sum_{j=1}^u K_j \xi_j^{(1,j)}; \rho_i^{(1,1)}, \dots, \rho_i^{(1,r)}, \tau_i^{(1,1)}, \dots, \tau_i^{(1,l)}, 1, 0, \dots, 0]_{1,s}, \dots,$$

$$[1 + \sigma_i^{(T)} - \eta_{G,g} c_i^{(T)} - \sum_{j=1}^u K_j \xi_j^{(T,j)}; \rho_i^{(T,1)}, \dots, \rho_i^{(T,r)}, \tau_i^{(T,1)}, \dots, \tau_i^{(T,l)}, 0, \dots, 0, 1]_{1,s},$$

$$[1 - A_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0]_{1,P},$$

$$[1 - \alpha_i - \eta_{G,g} a_i - \sum_{j=1}^u K_j \alpha_i^{(j)}; \delta'_i, \dots, \delta_i^{(r)}, \mu'_i, \dots, \mu_i^{(l)}, 1, \dots, 1, 0, \dots, 0]_{1,s},$$

W-items (T-W)-items

$$[1 - \beta_i - \eta_{G,g} b_i - \sum_{j=1}^u K_j \beta_i^{(j)}; \eta'_i, \dots, \eta_i^{(r)}, \theta'_i, \dots, \theta_i^{(l)}, 0, \dots, 0, 1, \dots, 1]_{1,s} \quad (4.3)$$

W-items (T-W)-items

$$B^* = [1 + \sigma'_i - \eta_{G,g} c'_i - \sum_{j=1}^u K_j \xi_i^{(1,j)}; \rho_i^{(1,1)}, \dots, \rho_i^{(1,r)}, \tau_i^{(1,1)}, \dots, \tau_i^{(1,l)}, 0, \dots, 0]_{1,s}, \dots,$$

$$[1 + \sigma_i^{(T)} - \eta_{G,g} c_i^{(T)} - \sum_{j=1}^u K_j \xi_i^{(T,j)}; \rho_i^{(T,1)}, \dots, \rho_i^{(T,r)}, \tau_i^{(T,1)}, \dots, \tau_i^{(T,l)}, 0, \dots, 0]_{1,s},$$

$$[1 - B_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0]_{1,Q},$$

$$[1 - \alpha_i - \beta_i - \eta_{G,g} (a_i + b_i) - \sum_{j=1}^u K_j (\alpha_i^{(j)} + \beta_i^{(j)}); (\delta'_i + \eta'_i), \dots, (\delta_i^{(r)} + \eta_i^{(r)})$$

$$(\mu'_i + \theta'_i), \dots, (\mu_i^{(l)} + \theta_i^{(l)}), 1, \dots, 1]_{1,s} \quad (4.4)$$

We have the following multiple Eulerian integral and we obtain the I-function of  $(r + l + T)$ -variables

$$\begin{aligned}
& \int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[ (x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right] \\
& \mathfrak{N}_{P_i, Q_i, c_i; r'}^{M, N} \left[ z \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{a_i} (v_i - x_i)^{b_i}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{c_i^{(j)}}} \right] \right] \\
& S_{N_1, \dots, N_u}^{M_1, \dots, M_u} \left( \begin{array}{c} y_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha'_i} (v_i - x_i)^{\beta'_i}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\xi_i^{(j,1)}}} \right] \\ \vdots \\ y_u \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha_i^{(u)}} (v_i - x_i)^{\beta_i^{(u)}}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\xi_i^{(j,u)}}} \right] \end{array} \right) \\
& \mathfrak{N}_{U; W}^{0, \mathbf{n}; V} \left( \begin{array}{c} z_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta'_i} (v_i - x_i)^{\eta'_i}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\rho_i^{(j,r)}}} \right] \end{array} \right) \\
& {}^P F_Q \left[ (A_P); (B_Q); - \sum_{k=1}^l g_k \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\mu_i^{(k)}} (v_i - x_i)^{\theta_i^{(r)}}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\tau_i^{(j,k)}}} \right] \right] dx_1 \cdots dx_s \\
& = \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{i=1}^s \left[ (v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{\sigma_i^{(j)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{\sigma_i^{(j)}} \right] \\
& \sum_{G=1}^M \sum_{g=0}^{\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_u=0}^{[N_u/M_u]} a_u \frac{(-)^g \Omega_{P_i, Q_i, c_i, r'}^{M, N}(\eta_{G, g})}{B_G g!} y_1^{K_1} \cdots y_u^{K_u} \prod_{i=1}^s \left[ (v_i - u_i)^{\eta_{G, g}(a_i + b_i) + \sum_{j=1}^u K_j (\alpha_i^{(j)} + \beta_i^{(j)})} \right]
\end{aligned}$$

$$z^{\eta_{G,g}} \prod_{i=1}^s \left[ \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\eta_{G,g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j,l)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\eta_{G,g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j,l)}} \right]$$

$$\mathfrak{N}_{U;sT+P+2s;V_1}^{0;n;sT+P+2s;V_1} \left( \begin{array}{c|c} \begin{matrix} z_1 w_1 \\ \vdots \\ z_r w_r \\ g_1 W_1 \\ \vdots \\ g_l W_l \\ G_1 \\ \vdots \\ G_T \end{matrix} & \begin{matrix} A ; A^* : C_1 \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ B ; B^* : D_1 \end{matrix} \end{array} \right) \quad (4.5)$$

Where

$$w_m = \prod_{i=1}^s \left[ (v_i - u_i)^{\delta_i^{(m)} + \eta_i^{(m)}} \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\rho_i^{(j,m)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\rho_i^{(j,m)}} \right], m = 1, \dots, r \quad (4.6)$$

$$W_k = \prod_{i=1}^s \left[ (v_i - u_i)^{\mu_i^{(k)} + \theta_i^{(k)}} \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\tau_i^{(j,k)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\tau_i^{(j,k)}} \right], k = 1, \dots, l \quad (4.7)$$

$$G_j = \prod_{i=1}^s \left[ \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = 1, \dots, W \quad (4.8)$$

$$G_j = - \prod_{i=1}^s \left[ \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = W + 1, \dots, T \quad (4.9)$$

Provided that :

$$\textbf{(A)} \ 0 \leq W \leq T; u_i, v_i \in \mathbb{R}; \min\{a_i, b_i, c_i^{(j)}, \alpha_i^{(k)}, \beta_i^{(k)}, \xi_i^{(j,k)}\} > 0, i = 1, \dots, s; k = 1, \dots, u; j = 1, \dots, T$$

$$\textbf{(B)} \ \min\{\delta_i^{(t)}, \eta_i^{(t)}, \rho_i^{(j,t)}, \mu_i^{(k)}, \theta_i^{(k)}, \tau_i^{(j,k)}\} \geq 0; j = 1, \dots, T; i = 1, \dots, s; k = 1, \dots, l; t = 1, \dots, r$$

$$\textbf{(C)} \ \sigma_i^{(j)} \in \mathbb{R}, U_i^{(j)}, V_i^{(j)} \in \mathbb{C}, z_t, g_k \in \mathbb{C}; j = 1, \dots, s; t = 1, \dots, r; k = 1, \dots, l; t = 1, \dots, r$$

$$\textbf{(D)} \ \max \left[ \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = 1, \dots, W \text{ and}$$

$$\max \left[ \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = W + 1, \dots, T$$

$$(F) \operatorname{Re} \left[ \alpha_i + a_i \min_{1 \leq j \leq M} \frac{b_j}{B_j} + \sum_{t=1}^r \delta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{d_j^{(t)}}{\delta_j^{(t)}} \right] > 0;$$

$$\operatorname{Re} \left[ \beta_i + b_i \min_{1 \leq j \leq M} \frac{b_j}{B_j} + \sum_{t=1}^r \eta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{d_j^{(t)}}{\delta_j^{(t)}} \right] > 0; i = 1, \dots, s$$

$$(G) U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j'^{(k)} \\ - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} \leq 0$$

$$(H) A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} \\ + \sum_{j=1}^{m_k} \delta_j'^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} - \delta_i^{(k)} - \eta_i^{(k)} - \sum_{j=1}^T \rho_i^{(j,k)} > 0$$

with  $k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$

$$(I) \left| \arg \left( z_i \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\rho_i^{(j,k)}} \right) \right| < \frac{1}{2} A_i^{(k)} \pi$$

(H) See I

(J)  $P \leq Q + 1$ . The equality holds, when , in addition,

$$\text{either } P > Q \text{ and } \sum_{k=1}^l \left| g_k \left( \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\tau_i^{(j,k)}} \right) \right|^{\frac{1}{Q-P}} < 1 \quad (u_i \leq x_i \leq v_i; i = 1, \dots, s)$$

$$\text{or } P \leq Q \text{ and } \max_{1 \leq k \leq l} \left[ \left| \left( g_k \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\tau_i^{(j,k)}} \right) \right| \right] < 1 \quad (u_i \leq x_i \leq v_i; i = 1, \dots, s)$$

(K) The series occuring on the right-hand side of (4.5) are absolutely and uniformly convergent

**Proof**

To establish the formula (4.5), we first use series representation (1.15) and (1.17) for  $\aleph_{P_i, Q_i, c_i; r}^{M, N}(\cdot)$  and  $S_{N_1, \dots, N_u}^{M_1, \dots, M_u}[\cdot]$  respectively, we use contour integral representation with the help of (1.1) for the multivariable Aleph-function occurring in its left-hand side and use the contour integral representation with the help of (3.1) for the generalized hypergeometric function  ${}_P F_Q(\cdot)$ . Changing the order of integration and summation (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process). Now we write :

$$\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}}$$

$$\text{where } K_i^{(j)} = \eta_i^{(j)} - \eta_{G,g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j,l)} - \sum_{t=1}^r \rho_i^{(j,t)} \zeta_t - \sum_{k=1}^l \tau_i^{(j,k)} \zeta_k \quad i = 1, \dots, s; j = 1, \dots, T \quad (4.10)$$

and express the factor occuring in R.H.S. Of (4.5) in terms of following Mellin-Barnes contour integral with the help of the result [8, page 18, eq.(2.6.4)]

$$\frac{1}{(2\pi\omega)^W} \int_{L'_1} \cdots \int_{L'_W} \prod_{j=1}^W [\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j)] \prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W \left[ \frac{(U_i^{(j)} u_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \prod_{j=1}^W \left[ \frac{(U_i^{(j)} (x_i - u_i))^{\zeta'_j}}{(u_i U_i^{(j)} + V_i^{(j)})^{\zeta'_j}} \right] d\zeta'_1 \cdots d\zeta'_W \quad (4.11)$$

and

$$\frac{1}{(2\pi\omega)^{T-W}} \int_{L'_{W+1}} \cdots \int_{L'_{Tj=W+1}} \prod_{j=W+1}^T [\Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j)] \prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=W+1}^T \left[ \frac{(U_i^{(j)} v_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \prod_{j=W+1}^T \left[ -\frac{(U_i^{(j)} (v_i - x_i))^{\zeta'_j}}{(v_i U_i^{(j)} + V_i^{(j)})^{\zeta'_j}} \right] d\zeta'_{W+1} \cdots d\zeta'_T \quad (4.12)$$

We apply the Fubini's theorem for multiple integral. Finally evaluating the innermost  $\mathbf{x}$ -integral with the help of (3.2) and reinterpreting the multiple Mellin-Barnes contour integral in terms of multivariable Aleph-function of  $(r + l + T)$ -variables, we obtain the formula (4.5).

## 5. Second formula

We note :

$$V_1 = V; 1, 1; 1, 0; \cdots; 1, 0; W_1 = W; 0, 1; \cdots; 0, 1 \quad (5.1)$$

$$C_1 = C; (1, 0); \cdots; (1, 0); D_1 = D; (0, 1), \cdots, (0, 1) \quad (5.2)$$

$$A^* = [1 + \sigma'_i - \theta'_i R - \eta_{G,g} c'_i - \sum_{j=1}^u K_j \xi_i^{(1,j)}; \rho_i^{(1,1)}, \cdots, \rho_i^{(1,r)}, 1, 0, \cdots, 0]_{1,s}, \cdots,$$

$$[1 + \sigma_i^{(T)} - \theta_i^{(T)} R - \eta_{G,g} c_i^{(T)} - \sum_{j=1}^u K_j \xi_i^{(T,j)}; \rho_i^{(T,1)}, \cdots, \rho_i^{(T,r)}, 0, \cdots, 0, 1]_{1,s},$$

$$[1 - \alpha_i - \zeta_i R - \eta_{G,g} a_i - \sum_{j=1}^u K_j \alpha_i^{(j)}; \delta'_i, \cdots, \delta_i^{(r)}, 1, \cdots, 1, 0, \cdots, 0]_{1,s},$$

W-items (T-W)-items

$$[1 - \beta_i - \lambda_i R - \eta_{G,g} b_i - \sum_{j=1}^u K_j \beta_i^{(j)}; \eta'_i, \cdots, \eta_i^{(r)}, 0, \cdots, 0, 1, \cdots, 1]_{1,s} \quad (5.3)$$

W-items (T-W)-items

$$B^* = [1 + \sigma'_i - \theta'_i R - \eta_{G,g} c'_i - \sum_{j=1}^u K_j \xi_i^{(1,j)}; \rho_i^{(1,1)}, \cdots, \rho_i^{(1,r)}, 0, \cdots, 0]_{1,s}, \cdots,$$

$$[1 + \sigma_i^{(T)} - \theta_i^{(T)} R - \eta_{G,g} c_i^{(T)} - \sum_{j=1}^u K_j \xi_i^{(T,j)}; \rho_i^{(T,1)}, \dots, \rho_i^{(T,r)}, 0, \dots, 0]_{1,s},$$

$$[1 - \alpha_i - \beta_i (\zeta_i + \lambda_i) R - \eta_{G,g} (a_i + b_i) - \sum_{j=1}^u K_j (\alpha_i^{(j)} + \beta_i^{(j)}); (\delta'_i + \eta'_i), \dots, (\delta_i^{(r)} + \eta_i^{(r)})$$

$$(\mu'_i + \theta'_i), \dots, (\mu_i^{(l)} + \theta_i^{(l)}), 1, \dots, 1]_{1,s} \quad (5.4)$$

We have the following multiple Eulerian integral

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[ (x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\mathfrak{N}_{P_i, Q_i, c_i; r'}^{M, N} \left[ z \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha_i} (v_i - x_i)^{\beta_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{c_i^{(j)}}} \right] \right] R_n^{\alpha, \beta} \left[ Z \prod_{j=1}^s \left[ \frac{(x_i - u_i)^{\zeta_i} (v_i - x_i)^{\lambda_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\theta_i^{(j)}}} \right] \right]$$

$$S_{N_1, \dots, N_u}^{M_1, \dots, M_u} \left( \begin{array}{c} y_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha'_i} (v_i - x_i)^{\beta'_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,1)}}} \right] \\ \vdots \\ y_u \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha_i^{(u)}} (v_i - x_i)^{\beta_i^{(u)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,u)}}} \right] \end{array} \right)$$

$$\mathfrak{N}_{U:W}^{0, n:V} \left( \begin{array}{c} z_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta'_i} (v_i - x_i)^{\eta'_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{array} \right) dx_1 \cdots dx_s$$

$$= \prod_{i=1}^s \left[ (v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\sum_{G=1}^M \sum_{g=0}^{\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_u=0}^{[N_u/M_u]} a_u \frac{(-)^g \Omega_{P_i, Q_i, c_i, r'}^{M, N} (\eta_{G,g})}{B_G g!} y_1^{K_1} \cdots y_u^{K_u}$$

$$\sum_{w,v,u,t,e,k_1,k_2} \psi'(w, v, u, t, e, k_1, k_2) Z^R \prod_{i=1}^s \left[ (v_i - u_i)^{\eta_{G,g}(a_i+b_i) + \sum_{j=1}^u K_j(\alpha_i^{(j)} + \beta_i^{(j)})} \right]$$

$$\prod_{i=1}^s \left[ \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\eta_{G,g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j,l)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\eta_{G,g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j,l)}} \right]$$

$$\mathbb{N}_{U;sT+2s;sT+s:W_1}^{0;n;sT+2s:V_1} \left( \begin{array}{c|c} \mathbf{z}_1 w_1 & \mathbf{A}; \mathbf{A}^* : C_1 \\ \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ \mathbf{z}_r w_r & \cdot \\ & \cdot \\ & \cdot \\ \mathbf{G}_1 & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ \mathbf{G}_T & \mathbf{B}; \mathbf{B}^* : D_1 \end{array} \right) \quad (5.5)$$

where

$$\psi'(w, v, u, t, e, k_1, k_2) = \frac{\psi(w, v, u, t, e, k_1, k_2, ) \prod_{i=1}^s (v_i - u_i)^{(\zeta_i + \lambda_i)R}}{\prod_{i=1}^s \left[ \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\theta_i^{(j)} R} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\theta_i^{(j)} R} \right]} \quad (5.6)$$

$\psi(w, v, u, t, e, k_1, k_2)$  and  $R$  are given by (2.4) and (2.3) respectively.

$$w_l = \prod_{i=1}^s \left[ (v_i - u_i)^{\delta_i^{(l)} + \eta_i^{(l)}} \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\rho_i^{(j,l)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\rho_i^{(j,l)}} \right], l = 1, \dots, r \quad (5.7)$$

$$G_j = \prod_{i=1}^s \left[ \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = 1, \dots, W \quad (5.8)$$

$$G_j = - \prod_{i=1}^s \left[ \frac{(v_i - u_i) U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right], j = W + 1, \dots, T \quad (5.9)$$

Provided that :

**(A)**  $0 \leq W \leq T; u_i, v_i \in \mathbb{R}; \min\{a_i, b_i, c_i^{(j)}, \alpha_i^{(k)}, \beta_i^{(k)}, \xi_i^{(j,k)}\} > 0, i = 1, \dots, s; k = 1, \dots, u; j = 1, \dots, T$

**(B)**  $\min\{\zeta_i, \lambda_i, \theta_i^{(j)}, \delta_i^{(l)}, \eta_i^{(l)}, \rho_i^{(j,l)}\} \geq 0; j = 1, \dots, T; i = 1, \dots, s; l = 1, \dots, r$

**(C)**  $\operatorname{Re}(\alpha_i, \beta_i, v_i^{(j)}) > 0 (i = 1, \dots, s; j = 1, \dots, T); |\tau| < 1$

$$(D) \max \left[ \frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = 1, \dots, W \text{ and}$$

$$\max \left[ \frac{(v_i - u_i)U_i^{(j)}}{u_i U_i^{(j)} + V_i^{(j)}} \right] < 1, i = 1, \dots, s; j = W + 1, \dots, T$$

$$(E) U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i(k)} \sum_{j=n_k+1}^{p_i(k)} \gamma_{ji(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} \\ - \tau_{i(k)} \sum_{j=m_k+1}^{q_i(k)} \delta_{ji(k)} \leq 0$$

$$(F) A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i(k)} \sum_{j=n_k+1}^{p_i(k)} \gamma_{ji(k)} \\ + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i(k)} \sum_{j=m_k+1}^{q_i(k)} \delta_{ji(k)} - \delta_i^{(k)} - \eta_i^{(k)} - \sum_{j=1}^T \rho_i^{(j,k)} > 0$$

$$(i = 1, \dots, s; k = 1, \dots, r)$$

$$(G) Re[\alpha_i + R\zeta_i + a_i \min_{1 \leq j \leq M} \frac{b_j}{B_j} + \sum_{t=1}^r \delta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{d_j^{(t)}}{\delta_j^{(t)}}] > 0;$$

$$Re[\beta_i + R\lambda_i + b_i \min_{1 \leq j \leq M} \frac{b_j}{B_j} + \sum_{t=1}^r \eta_i^{(t)} \min_{1 \leq j \leq m_i} \frac{d_j^{(t)}}{\delta_j^{(t)}}] > 0; i = 1, \dots, s$$

$$(H) \left| arg \left( z_i \prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{-\rho_i^{(j,k)}} \right) \right| < \frac{1}{2} A_i^{(k)} \pi$$

$$(I) \text{ See I}$$

$$(J) \text{ The series occuring on the right-hand side of (5.5) are absolutely and uniformly convergent}$$

### Proof

To establish the formula (5.5), we first use series representation (1.15), (1.17) and (3.1) for  $\aleph_{P_i, Q_i, c_i; r}^{M, N}(\cdot)$ ,  $S_{N_1, \dots, N_u}^{M_1, \dots, M_u}[\cdot]$  and  $R_n^{\alpha, \beta}[\cdot]$  respectively and the contour integral representation with the help of (1.2) for the multivariable Aleph-function occuring in its left-hand side. Changing the order of integration and summation (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process). Now, we write:

$$\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} \quad (5.10)$$

$$\text{where } K_i^{(j)} = \eta_i^{(j)} - \eta_{G,g} c_i^{(j)} - R\theta_i^{(j)} - \sum_{l=1}^u L_l \xi_i^{(j,l)} - \sum_{t=1}^r \rho_i^{(j,t)} \zeta_t; i = 1, \dots, s; j = 1, \dots, T \quad (5.11)$$

and express the factors occuring in R.H.S. Of (5.5) in terms of following Mellin-Barnes contour integral , we obtain :

$$\prod_{j=1}^W (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=1}^W \left[ \frac{(U_i^{(j)} u_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^W} \int_{L'_1} \cdots \int_{L'_W} \prod_{j=1}^W \left[ \Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right. \\ \left. \prod_{j=1}^W \left[ \frac{(U_i^{(j)}(x_i - u_i))^{\zeta'_j}}{(u_i U_i^{(j)} + V_i^{(j)})} \right] \right] d\zeta'_1 \cdots d\zeta'_W \quad (5.12)$$

and

$$\prod_{j=W+1}^T (U_i^{(j)} x_i + V_i^{(j)})^{K_i^{(j)}} = \prod_{j=W+1}^T \left[ \frac{(U_i^{(j)} v_i + V_i^{(j)})^{K_i^{(j)}}}{\Gamma(-K_i^{(j)})} \right] \frac{1}{(2\pi\omega)^{T-W}} \int_{L'_{W+1}} \cdots \int_{L'_T} \prod_{j=W+1}^T \left[ \Gamma(-\zeta'_j) \Gamma(-K_i^{(j)} + \zeta'_j) \right. \\ \left. \prod_{j=W+1}^T \left[ \frac{(U_i^{(j)}(x_i - v_i))^{\zeta'_j}}{(v_i U_i^{(j)} + V_i^{(j)})} \right] \right] d\zeta'_{W+1} \cdots d\zeta'_T \quad (5.13)$$

We apply the Fubini's theorem for multiple integral. Finally evaluating the innermost  $\mathbf{x}$ -integral with the help of (3.2) and reinterpreting the multiple Mellin-Barnes contour integral in terms of multivariable Aleph-function of  $(r+T)$ -variables, we obtain the formula (5.5).

## 6. Multivariable I-function

If  $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$ , the multivariable Aleph-functions reduces to multivariable I-function defined by Sharma and al [ 6] and we have two general multiple Eulerian integrals of multivariable I-functions.

### Formula 1

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[ (x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\mathfrak{N}_{P_i, Q_i, c_i; r'}^{M, N} \left[ \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{a_i} (v_i - x_i)^{b_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{c_i^{(j)}}} \right] \right]$$

$$S_{N_1, \dots, N_u}^{M_1, \dots, M_u} \left( \begin{array}{c} y_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha'_i} (v_i - x_i)^{\beta'_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,1)}}} \right] \\ \vdots \\ y_u \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha_i^{(u)}} (v_i - x_i)^{\beta_i^{(u)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\xi_i^{(j,u)}}} \right] \end{array} \right)$$

$$I_{U:W}^{0,n;V} \left( \begin{array}{c} z_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta'_i} (v_i - x_i)^{\eta'_i}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\rho_i^{(j,r)}}} \right] \end{array} \right)$$

$${}_P F_Q \left[ (A_P); (B_Q); - \sum_{k=1}^l g_k \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\mu_i^{(k)}} (v_i - x_i)^{\theta_i^{(r)}}}{\prod_{j=1}^T (U_i^{(j)} x_i + V_i^{(j)})^{\tau_i^{(j,k)}}} \right] \right] dx_1 \cdots dx_s$$

$$= \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{i=1}^s \left[ (v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\sum_{G=1}^M \sum_{g=0}^{\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_u=0}^{[N_u/M_u]} a_u \frac{(-)^g \Omega_{P_i, Q_i, c_i, r'}^{M, N}(\eta_{G, g})}{B_G g!} y_1^{K_1} \cdots y_u^{K_u} \prod_{i=1}^s \left[ (v_i - u_i)^{\eta_{G, g}(a_i + b_i) + \sum_{j=1}^u K_j (\alpha_i^{(j)} + \beta_i^{(j)})} \right]$$

$$z^{\eta_{G,g}} \prod_{i=1}^s \left[ \prod_{j=1}^W (u_i U_i^{(j)} + V_i^{(j)})^{-\eta_{G,g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(l,k)}} \prod_{j=W+1}^T (u_i U_i^{(j)} + V_i^{(j)})^{-\eta_{G,g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j,l)}} \right]$$

$$I_{U;sT+P+2s;V_1}^{0;n;sT+P+2s;sT+Q+s;W_1} \left( \begin{array}{c|c} \begin{array}{c} z_1 w_1 \\ \vdots \\ z_r w_r \\ g_1 W_1 \\ \vdots \\ g_l W_l \\ G_1 \\ \vdots \\ G_T \end{array} & \begin{array}{c} A ; A^* : C_1 \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ B ; B^* : D_1 \end{array} \end{array} \right) \quad (6.1)$$

under the same conditions and notations that (4.5) with  $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$

**Formula 2**

$$\int_{u_1}^{v_1} \cdots \int_{u_s}^{v_s} \prod_{i=1}^s \left[ (x_i - u_i)^{\alpha_i - 1} (v_i - x_i)^{\beta_i - 1} \prod_{j=1}^T (u_i^{(j)} x_i + v_i^{(j)})^{\sigma_i^{(j)}} \right]$$

$$\mathfrak{N}_{P_i, Q_i, c_i; r'}^{M, N} \left[ z \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{a_i} (v_i - x_i)^{b_i}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{c_i^{(j)}}} \right] \right] R_n^{\alpha, \beta} \left[ Z \prod_{j=1}^s \left[ \frac{(x_i - u_i)^{\zeta_i} (v_i - x_i)^{\lambda_i}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\theta_i^{(j)}}} \right] \right]$$

$$S_{N_1, \dots, N_u}^{M_1, \dots, M_u} \left( \begin{array}{c} y_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha'_i} (v_i - x_i)^{\beta'_i}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\xi_i^{(j,1)}}} \right] \\ \vdots \\ y_u \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\alpha_i^{(u)}} (v_i - x_i)^{\beta_i^{(u)}}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\xi_i^{(j,u)}}} \right] \end{array} \right)$$

$$I_{U:W}^{0, \mathbf{n}; V} \left( \begin{array}{c} z_1 \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta'_i} (v_i - x_i)^{\eta'_i}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\rho_i^{(j,1)}}} \right] \\ \vdots \\ z_r \prod_{i=1}^s \left[ \frac{(x_i - u_i)^{\delta_i^{(r)}} (v_i - x_i)^{\eta_i^{(r)}}}{\prod_{j=1}^T \left( U_i^{(j)} x_i + V_i^{(j)} \right)^{\rho_i^{(j,r)}}} \right] \end{array} \right) dx_1 \cdots dx_s$$

$$= \prod_{i=1}^s \left[ (v_i - u_i)^{\alpha_i + \beta_i - 1} \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{\sigma_i^{(j)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{\sigma_i^{(j)}} \right]$$

$$\sum_{G=1}^M \sum_{g=0}^{\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_u=0}^{[N_u/M_u]} a_u \frac{(-)^g \Omega_{P_i, Q_i, c_i, r'}^{M, N}(\eta_{G, g})}{B_G g!} y_1^{K_1} \cdots y_u^{K_u}$$

$$\sum_{w, v, u, t, e, k_1, k_2} \psi'(w, v, u, t, e, k_1, k_2) Z^R \prod_{i=1}^s \left[ (v_i - u_i)^{\eta_{G, g} (a_i + b_i) + \sum_{j=1}^u K_j (\alpha_i^{(j)} + \beta_i^{(j)})} \right]$$

$$\prod_{i=1}^s \left[ \prod_{j=1}^W \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\eta_{G, g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j, l)}} \prod_{j=W+1}^T \left( u_i U_i^{(j)} + V_i^{(j)} \right)^{-\eta_{G, g} c_i^{(j)} - \sum_{l=1}^u K_l \xi_i^{(j, l)}} \right]$$

$$I_{U;sT+2s;sT+s;W_1}^{0;n;sT+2s;V_1} \left( \begin{array}{c|c} z_1 w_1 & A ; A^* : C_1 \\ \cdot & \cdot \\ \cdot & \cdot \\ z_r w_r & \cdot \\ & \cdot \\ G_1 & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ G_T & B ; B^* : D_1 \end{array} \right) \quad (6.2)$$

under the same conditions and notations that (5.5) with  $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$

## 9. Conclusion

In this paper we have evaluated two generalized multiple Eulerian integrals involving the multivariable Aleph-functions and multivariable I-function defined by Sharma et [6] with general arguments. The formulae established in this paper is very general nature. Thus, the results established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

## REFERENCES

- [1] Agrawal B.D. And Chaubey J.P. Certain derivation of generating relations for generalized polynomials. Indian J. Pure and Appl. Math 10 (1980), page 1155-1157, ibid 11 (1981), page 357-359
- [2] Chaurasia V.B.L and Singh Y. New generalization of integral equations of fredholm type using Aleph-function Int. J. of Modern Math. Sci. 9(3), 2014, p 208-220.
- [3] Fujiwara I. A unified presentation of classical orthogonal polynomials. Math. Japon. 11 (1966), page133-148.
- [4] Raizada S.K. A study of unified representation of special functions of Mathematics Physics and their use in statistical and boundary value problem. Ph.D. Thesis, Bundelkhand University, Jhansi, India, 1991
- [5] Salim T.O. A serie formula of generalized class of polynomials associated with Laplace transform and fractional integral operators. J. Rajasthan. Acad. Phy. Sci. 1(3) (2002), page 167-176.
- [6] Sharma C.K.and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20,no2, p 113-116.
- [7] Srivastava H.M. A multilinear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial, Pacific. J. Math. 177(1985), page183-191.
- [8] Srivastava H.M. Gupta K.C. And Goyal S.P. The H-functions of one and two variables with applications. South Asian Publishers Pvt Ltd 1982
- [9] Srivastava H.M. and Karlsson P.W. Multiple Gaussian Hypergeometric series. Ellis.Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.
- [10] Südland N.; Baumann, B. and Nonnenmacher T.F. , Open problem : who knows about the Aleph-functions? Fract. Calc. Appl. Anal., 1(4) (1998): 401-402.

