Fractional derivatives involving general classes of polynomials and special

functions representable by Mellin-Barnes type contour integrals III

$F.Y. AYANT^1$

1 Teacher in High School , France E-mail : fredericayant@gmail.com

ABSTRACT

In this document, first time, we established fractional derivative involving the product of two general classes of polynomials with Multivariable Ifunction defined by Nambisan et al [6] and generalized hypergeometric function. Second time, we established fractional derivative involving the product of two general classes of polynomials with multivariable I-functions and generalized hypergeometric function.

Keywords: Multivariable Aleph-function, general class of polynomial, fractional derivative, generalized hypergeometric function, multivariable I-function, multivariable H-function.

2010 Mathematics Subject Classification. 33C45, 33C60, 26D20

1.Introduction and preliminaries.

In this document, first time, we established fractional derivative involving the product of two general classes of polynomials with Multivariable I-function defined by Nambisan et al [6] and generalized hypergeometric function. Second time, we established fractional derivative involving the product of two general classes of polynomials with multivariable I-functions and generalized hypergeometric function.

The I-function is defined and represented in the following manner.

$$I(z_1, z_2, ... z_r) = I_{p,q:p_1,q_1; \cdots; p_r,q_r}^{0,n:m_1,n_1; \cdots; m_r,n_r} \begin{pmatrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{pmatrix} (a_j; \alpha_j^{(1)}, \cdots, \alpha_j^{(r)}; A_j)_{1,p} :$$

$$(c_{j}^{(1)}, \gamma_{j}^{(1)}; C_{j}^{(1)})_{1,p_{1}}; \cdots; (c_{j}^{(r)}, \gamma_{j}^{(r)}; C_{j}^{(r)})_{1,p_{r}}$$

$$(d_{j}^{(1)}, \delta_{j}^{(1)}; D_{j}^{(1)})_{1,q_{1}}; \cdots; (d_{j}^{(r)}, \delta_{j}^{(r)}; D_{j}^{(r)})_{1,q_{r}}$$

$$(1.1)$$

$$=\frac{1}{(2\pi\omega)^r}\int_{L_1}\cdots\int_{L_r}\phi(s_1,\cdots,s_r)\prod_{i=1}^r\theta_i(s_i)z_i^{s_i}\mathrm{d}s_1\cdots\mathrm{d}s_r$$
(1.2)

where $\phi(s_1,\cdots,s_r)$, $heta_i(s_i)$, $i=1,\cdots,r$ are given by :

$$\phi(s_1, \cdots, s_r) = \frac{\prod_{j=1}^n \Gamma^{A_j} \left(1 - a_j + \sum_{i=1}^r \alpha_j^{(i)} s_j \right)}{\prod_{j=n+1}^p \Gamma^{A_j} \left(a_j - \sum_{i=1}^r \alpha_j^{(i)} s_j \right) \prod_{j=1}^q \Gamma^{B_j} \left(1 - b_j + \sum_{i=1}^r \beta_j^{(i)} s_j \right)}$$
(1.3)

$$\theta_{i}(s_{i}) = \frac{\prod_{j=1}^{n_{i}} \Gamma^{C_{j}^{(i)}} \left(1 - c_{j}^{(i)} + \gamma_{j}^{(i)} s_{i}\right) \prod_{j=1}^{m_{i}} \Gamma^{D_{j}^{(i)}} \left(d_{j}^{(i)} - \delta_{j}^{(i)} s_{i}\right)}{\prod_{j=n_{i}+1}^{p_{i}} \Gamma^{C_{j}^{(i)}} \left(c_{j}^{(i)} - \gamma_{j}^{(i)} s_{i}\right) \prod_{j=m_{i}+1}^{q_{i}} \Gamma^{D_{j}^{(i)}} \left(1 - d_{j}^{(i)} - \delta_{j}^{(i)} s_{i}\right)} \right)}$$
(1.4)

For more details, see Nambisan et al [6].

Following the result of Braaksma [1] the I-function of r variables is analytic if :

$$U_{i} = \sum_{j=1}^{p} A_{j} \alpha_{j}^{(i)} - \sum_{j=1}^{q} B_{j} \beta_{j}^{(i)} + \sum_{j=1}^{p_{i}} C_{j}^{(i)} \gamma_{j}^{(i)} - \sum_{j=1}^{q_{i}} D_{j}^{(i)} \delta_{j}^{(i)} \leqslant 0, i = 1, \cdots, r$$
(1.5)

The integral (2.1) converges absolutely if

$$|arg(z_k)| < \frac{1}{2} \Delta_k \pi, k = 1, \cdots, r \text{ where}$$

$$\Delta_k = -\sum_{j=n+1}^p A_j \alpha_j^{(k)} - \sum_{j=1}^q B_j \beta_j^{(k)} + \sum_{j=1}^{m_k} D_j^{(k)} \delta_j^{(k)} - \sum_{j=m_k+1}^{q_k} D_j^{(k)} \delta_j^{(k)} + \sum_{j=1}^{n_k} C_j^{(k)} \gamma_j^{(k)} - \sum_{j=n_k+1}^{p_k} C_j^{(k)} \gamma_j^{(k)} > 0 \quad (1.6)$$

Consider the second multivariable I-function.

$$(\mathbf{c}_{j}^{(1)}, \gamma_{j}^{\prime(1)}; C_{j}^{\prime(1)})_{1, p_{1}^{\prime}}; \cdots; (c_{j}^{\prime(s)}, \gamma_{j}^{\prime(s)}; C_{j}^{\prime(s)})_{1, p_{s}^{\prime}})$$

$$(\mathbf{d}_{j}^{(1)}, \delta_{j}^{\prime(1)}; D_{j}^{\prime(1)})_{1, q_{1}^{\prime}}; \cdots; (d_{j}^{\prime(s)}, \delta_{j}^{\prime(s)}; D_{j}^{\prime(s)})_{1, q_{s}^{\prime}})$$

$$(1.7)$$

$$=\frac{1}{(2\pi\omega)^s}\int_{L_1}\cdots\int_{L_s}\psi(t_1,\cdots,t_s)\prod_{i=1}^s\xi_i(t_i)z_i^{t_i}\mathrm{d}t_1\cdots\mathrm{d}t_s$$
(1.8)

where $\ \psi(t_1,\cdots,t_s)$, $\xi_i(s_i)$, $i=1,\cdots,s$ are given by :

$$\psi(t_1, \cdots, t_s) = \frac{\prod_{j=1}^{n'} \Gamma^{A'_j} \left(1 - a'_j + \sum_{i=1}^s \alpha'_j{}^{(i)} t_j\right)}{\prod_{j=n'+1}^{p'} \Gamma^{A'_j} \left(a'_j - \sum_{i=1}^s \alpha'_j{}^{(i)} t_j\right) \prod_{j=1}^{q'} \Gamma^{B'_j} \left(1 - b'_j + \sum_{i=1}^s \beta'_j{}^{(i)} t_j\right)}$$
(1.9)

$$\xi_{i}(s_{i}) = \frac{\prod_{j=1}^{n'_{i}} \Gamma^{C'_{j}^{(i)}} \left(1 - c'_{j}^{(i)} + \gamma'_{j}^{(i)}t_{i}\right) \prod_{j=1}^{m'_{i}} \Gamma^{D'_{j}^{(i)}} \left(d'_{j}^{(i)} - \delta'_{j}^{(i)}t_{i}\right)}{\prod_{j=n'_{i}+1}^{p'_{i}} \Gamma^{C'_{j}^{(i)}} \left(c'_{j}^{(i)} - \gamma'_{j}^{(i)}t_{i}\right) \prod_{j=m'_{i}+1}^{q'_{i}} \Gamma^{D'_{j}^{(i)}} \left(1 - d'_{j}^{(i)} - \delta'_{j}^{(i)}t_{i}\right)}$$
(1.10)

For more details, see Nambisan et al [6].

Following the result of Braaksma [1] the I-function of r variables is analytic if :

$$U'_{i} = \sum_{j=1}^{p'} A'_{j} \alpha'_{j}{}^{(i)} - \sum_{j=1}^{q'} B'_{j} \beta'_{j}{}^{(i)} + \sum_{j=1}^{p'_{i}} C'_{j}{}^{(i)} \gamma'_{j}{}^{(i)} - \sum_{j=1}^{q'_{i}} D'_{j}{}^{(i)} \delta'_{j}{}^{(i)} \leqslant 0, i = 1, \cdots, s$$
(1.11)

The integral (2.1) converges absolutely if

where
$$|arg(z'_k)| < \frac{1}{2}\Delta'_k \pi, k = 1, \cdots, s$$

$$\Delta'_k = -\sum_{j=n'+1}^{p'} A'_j \alpha'_j{}^{(k)} - \sum_{j=1}^{q'} B'_j \beta'_j{}^{(k)} + \sum_{j=1}^{m'_k} D'^{(k)}_j \delta'_j{}^{(k)} - \sum_{j=m'_k+1}^{q'_k} D'^{(k)}_j \delta'_j{}^{(k)} + \sum_{j=1}^{n'_k} C'^{(k)}_j \gamma'_j{}^{(k)} - \sum_{j=n'_k+1}^{p'_k} C'^{(k)}_j \gamma'_j{}^{(k)} > 0$$
(1.12)

A general of Humbert polynomial is defined by the generating function [5,page 54, eq.(1.5)]

$$\left[C - axt + bt^m (2x - 1)^d\right]^v = \sum_{n=0}^{\infty} \phi_n(x) t^n$$
(1.13)

where $m, a, d \in N$, and other parameters are unrestricted in general.

The finite series representation for $\phi_n(x)$ is also obtained there [5,page 55 31. eq(2.1)]

$$\phi_n(x) = \sum_{k=0}^{[n/m]} \frac{(-)^k c^{-\nu - n + (m-1)k} (\nu)_{n+(1-m)k} (ax)^{n-mk} \left[b(2x-1)^d \right]^k}{k! (n-mk)!}$$
(1.14)

$$=\sum_{k=0}^{[n/m]}\sum_{l=0}^{[dk]}\frac{(-dk)_l 2^l c^{-\nu-n+(m-1)k}(\nu)_{n+(1-m)k}(a)^{n-mk}b^k(-)^{(d+1)k}x^{n-mk+l}}{l!k!(n-mk)!}$$
(1.15)

The generalized polynomials defined by Srivastava [8], is given in the following manner :

$$S_{N_{1}',\cdots,N_{t}'}^{M_{1}',\cdots,M_{t}'}[y_{1},\cdots,y_{t}] = \sum_{K_{1}=0}^{[N_{1}'/M_{1}']} \cdots \sum_{K_{t}=0}^{[N_{t}'/M_{t}']} \frac{(-N_{1}')_{M_{1}'K_{1}}}{K_{1}!} \cdots \frac{(-N_{t}')_{M_{t}'K_{t}}}{K_{t}!}$$

$$A[N_{1}',K_{1};\cdots;N_{t}',K_{t}]y_{1}^{K_{1}}\cdots y_{t}^{K_{t}}$$
(1.16)

Where M'_1, \dots, M'_s are arbitrary positive integers and the coefficients $A[N'_1, K_1; \dots; N'_t, K_t]$ are arbitrary constants, real or complex.

The another general class of polynomials of K-variables defined by Srivastava et al [9,page 686,eq.(1.4)] and denoted as follows

$$T_n^{m_1,\cdots,m_k}[x_1,\cdots,x_k] = \sum_{h_1,\cdots,h_k=0}^{M \leqslant n} \left[(-n)_M B(n,h_1,\cdots,h_k) \prod_{i=1}^k \frac{x_i^{h_i}}{h_i!} \right]$$
(1.17)

The Riemann-Liouville fractional derivative (or integral) of order μ is defined as follows [3,page49]

$$D_{x}^{\mu}f(x) = \begin{cases} \frac{1}{\Gamma(-u)} \int_{0}^{x} (x-t)^{-\mu-1} f(t) dt, Re(\mu) < 0 \\ & \ddots \\ & \\ \frac{d^{m}}{dx^{m}} [D_{x}^{\mu-m} \{f(x)\}], 0 \leqslant m, m \in \mathbb{N} \end{cases}$$
(1.18)

We have the following fractional derivative formula [4,page 67, eq. (4.4.4)] is also required :

$$D_x^{\mu}\left(x^{\lambda}\right) = \frac{\Gamma(\lambda+1)}{\Gamma(\lambda-\mu+1)} x^{\lambda-\mu}, Re(\lambda) > -1$$
(1.19)

The generalized Leibnitz formula for fractional calculus is required in the following form [7, page317]

$$D_x^{\alpha}[f(x)g(x)] = \sum_{n=-\infty}^{\infty} a\binom{\alpha}{an+\epsilon} D_x^{\alpha-an-\epsilon}[f(x)] D_x^{an+\epsilon}[g(x)]$$
(1.20)

 $0\leqslant a\leqslant 1;\alpha$ is a real or complex arbitrary number.

The following generalized hypergeometric function in terms of multiple contour integrals is also required [10,page 39 eq.30]

$$\frac{\prod_{j=1}^{P} \Gamma(A_j)}{\prod_{j=1}^{Q} \Gamma(B_j)} {}_{P}F_Q\left[(A_P); (B_Q); -(x_1 + \dots + x_t)\right]$$

$$= \frac{1}{(2\pi\omega)^{t}} \int_{L_{1}} \cdots \int_{L_{t}} \frac{\prod_{j=1}^{P} \Gamma(A_{j} + s_{1} + \dots + s_{t})}{\prod_{j=1}^{Q} \Gamma(B_{j} + s_{1} + \dots + s_{t})} \Gamma(-s_{1}) \cdots \Gamma(-s_{t}) x_{1}^{s_{1}} \cdots x_{t}^{s_{t}} \mathrm{d}s_{1} \cdots \mathrm{d}s_{t}$$
(1.21)

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of $\Gamma(A_j + s_1 + \cdots + s_t)$ are separated from those of $\Gamma(-s_j)$, $j = 1, \cdots, t$. The above result (1.21) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of $\Gamma(-s_j)$, $j = 1, \cdots, t$

2. Required formula

The fractional derivative of multivariable I-function is as follows :

Lemme

$$D_{z}^{\alpha}\left[z^{\lambda}I(c_{1}z^{\sigma_{1}},\cdots,c_{r}z^{\sigma_{r}})\right] = z^{\lambda-\alpha}I_{p+1,q:p_{1},q_{1};\cdots;p_{r},q_{r}}^{0,n+1:m_{1},n_{1};\cdots;m_{r},n_{r}} \begin{pmatrix} z_{1} \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ z_{r} \end{pmatrix} (a_{j};\alpha_{j}^{(1)},\cdots,\alpha_{j}^{(r)};A_{j})_{1,p},$$

$$(-\lambda; \sigma_{1}, \cdots, \sigma_{r}; 1): (c_{j}^{(1)}, \gamma_{j}^{(1)}; C_{j}^{(1)})_{1,p_{1}}; \cdots; (c_{j}^{(r)}, \gamma_{j}^{(r)}; C_{j}^{(r)})_{1,p_{r}}$$

$$(\alpha - \lambda; \sigma_{1}, \cdots, \sigma_{r}; 1): (d_{j}^{(1)}, \delta_{j}^{(1)}; D_{j}^{(1)})_{1,q_{1}}; \cdots; (d_{j}^{(r)}, \delta_{j}^{(r)}; D_{j}^{(r)})_{1,q_{r}}$$

$$(2.1)$$

To prove the Lemme ,express the multivariable I-function in Mellin-Barnes contour integral with the help of (1.2) and use the formula (1.19). Now interpreting the result thus obtained with the Mellin-barnes contour integral, we arrive at the desired result.

Where :
$$\sigma_i > 0, i = 1, \cdots, r$$
; $Re[\lambda + \sum_{i=1}^r \sigma_i \min_{1 \le j \le m_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + 1] > 0$

3. Main results

We will use these notations for the two following theorems:

$$X = m_1, n_1; \dots; m_r, n_r; 1, 0; \dots; 1, 0; V = p_1, q_1; \dots; p_r, q_r; 0, 1; \dots; 0, 1$$
(3.1)

$$A = (a_j; \alpha_j^{(1)}, \cdots, \alpha_j^{(r)}, 0, \cdots, 0; A_j)_{1,p}$$
(3.2)

$$B = (b_j; \beta_j^{(1)}, \cdots, \beta_j^{(r)}, 0, \cdots, 0; B_j)_{1,q}$$
(3.3)

$$\mathbf{C} = (\mathbf{c}_{j}^{(1)}, \gamma_{j}^{(1)}; C_{j}^{(1)})_{1,p_{1}}; \cdots; (c_{j}^{(r)}, \gamma_{j}^{(r)}; C_{j}^{(r)})_{1,p_{r}}; (1,0;1); \cdots; (1,0;1)$$
(3.4)

$$D = (\mathbf{d}_{j}^{(1)}, \delta_{j}^{(1)}; D_{j}^{(1)})_{1,q_{1}}; \cdots; (\mathbf{d}_{j}^{(r)}, \delta_{j}^{(r)}; D_{j}^{(r)})_{1,q_{r}}; (0, 1; 1); \cdots; (0, 1; 1)$$
(3.5)

$$K_1 = [1 - A_j; 0, \cdots, 0, 1, \cdots, 1; 1]_{1,P}, (1 + \lambda + \sum_{j=1}^k \delta_j h_j - l_1; \sigma_1, \cdots, \sigma_r, \rho'_1, \cdots, \rho'_t; 1),$$

$$(1 + \gamma + \sum_{j=1}^{k} \epsilon_{j} h_{j} - l_{2}; \mu_{1}, \cdots, \mu_{r}, \lambda_{1}', \cdots, \lambda_{t}'; 1),$$

$$(Mh - \mu - N - l - \upsilon l_{1} - \upsilon l_{2} - \sum_{j=1}^{k} \lambda_{j} h_{j}; \rho_{1}, \cdots, \rho_{r}, \delta_{1}', \cdots, \delta_{t}'; 1)$$

$$(3.6)$$

$$K_2 = [1 - B_j; 0, \cdots, 0, 1, \cdots, 1; 1]_{1,Q}, (1 + \lambda + \sum_{j=1}^n \delta_j h_j; \sigma_1, \cdots, \sigma_r, \rho'_1, \cdots, \rho'_t; 1),$$

$$(1 + \gamma + \sum_{j=1}^{k} \epsilon_{j} h_{j}; \mu_{1}, \cdots, \mu_{r}, \lambda_{1}', \cdots, \lambda_{t}'; 1),$$

$$(Mh - \mu + \alpha - N - l - \upsilon l_{1} - \upsilon l_{2} - \sum_{j=1}^{k} \lambda_{j} h_{j}; \rho_{1}, \cdots, \rho_{r}, \delta_{1}', \cdots, \delta_{t}'; 1)$$
(3.7)

$$D_x^{\alpha} \Big\{ x^{\mu} (x^{\upsilon} + \epsilon)^{\lambda} (x^{\upsilon} + \eta)^{\gamma} \phi_N(zx) S_{N_1', \cdots, N_k'}^{M_1', \cdots, M_k'} \begin{pmatrix} y_1 x^{\lambda_1} (x^{\upsilon} + \epsilon)^{\delta_1} (x^{\upsilon} + \eta)^{\epsilon_1} \\ \ddots \\ y_k x^{\lambda_k} (x^{\upsilon} + \epsilon)^{\delta_k} (x^{\upsilon} + \eta)^{\epsilon_k} \end{pmatrix}$$

$$I\left(\begin{array}{c} z_{1}x^{\rho_{1}}(x^{\upsilon}+\epsilon)^{-\sigma_{1}}(x^{\upsilon}+\eta)^{-\mu_{1}}\\ & \ddots\\ & & \\ z_{r}x^{\rho_{r}}(x^{\upsilon}+\epsilon)^{-\sigma_{r}}(x^{\upsilon}+\eta)^{-\mu_{r}} \end{array}\right)$$

$${}_{P}F_{Q}\left[(A_{P});(B_{Q});-\sum_{j=1}^{t}z_{j}'x^{\delta_{j}'}(x^{\upsilon}+\epsilon)^{-\rho_{j}'}(x^{\upsilon}+\eta)^{-\lambda_{j}'}\right]\right\}$$

$$=\frac{\epsilon^{\lambda}\eta^{\gamma}\prod_{j=1}^{Q}\Gamma(B_{j})}{\prod_{j=1}^{P}\Gamma(A_{j})}\sum_{h=0}^{[N/M]}\sum_{l=0}^{[dh]}\frac{(-dh)_{l}c^{-\upsilon'-N+(M-1)h}(\upsilon')_{N+(1-M)h}(a)^{N-Mh}b^{h}(-)^{(d+1)h}2^{l}x^{N-Mh+1}}{l!h!(N-Mh)!}$$

$$\sum_{h_{1}=0}^{[N_{1}'/M_{1}']} \cdots \sum_{h_{k}=0}^{[N_{k}'/M_{k}']} A(h_{1}, \cdots, h_{k}) \prod_{j=1}^{k} \left[\frac{(-N_{j}')_{M_{j}'h_{j}}}{h_{j}!} (x^{\lambda_{j}} \epsilon^{\delta_{j}} \eta^{\epsilon_{j}} y_{j})^{h_{j}} \right] \sum_{l_{1}, l_{2}=0}^{\infty} \frac{(-x)^{\upsilon l_{1}+\upsilon l_{2}}}{\epsilon^{l_{1}} l_{1}! \eta^{l_{2}} l_{2}!}$$

$$I_{0,n+P+3;X} = \left[\begin{pmatrix} z_{1} \epsilon^{-\sigma_{1}} \eta^{-\mu_{1}} x^{\rho_{1}} \\ \vdots \\ z_{r} \epsilon^{-\sigma_{r}} \eta^{-\mu_{r}} x^{\rho_{r}} \\ \vdots \\ z_{r} \epsilon^{-\sigma_{r}} \eta^{-\mu_{r}} x^{\rho_{r}} \\ \vdots \\ z_{r} \epsilon^{-\sigma_{r}} \eta^{-\mu_{r}} x^{\rho_{r}} \\ z_{r} t^{-\sigma_{r}} t^{-\sigma_{r}} x^{\rho_{r}} \\ z_{r} t^{-\sigma_{r}} t^{-\sigma_{r}} t^{-\sigma_{r}} t^{-\sigma_{r}} t^{-\sigma_{r}} t^{-\sigma_{r}} \\ z_{r} t^{-\sigma_{r}} t^{-\sigma$$

$$\begin{bmatrix}
 I_{p+P+3;X}^{0,n+P+3;X} \\
 z'_{1}\epsilon^{-\rho'_{1}}\eta^{-\lambda'_{1}}x^{\delta'_{1}} \\
 \vdots \\
 z'_{t}\epsilon^{-\rho'_{t}}\eta^{-\lambda'_{t}}x^{\delta'_{t}}
 \end{bmatrix}
 B, K_{2}:D$$
(3.10)

The quantities A, B, K_1, K_2, C and D are defined above, where the I-function on the right hand side is of (r+t) variables.

Provided that

a) $v, \lambda_i, \delta_i, \epsilon_i, \rho_j, \sigma_j, \mu_j, \delta'_p, \rho'_p, \lambda'_p > 0; i = 1, \cdots, k; j = 1, \cdots, r; p = 1, \cdots, t$

b)
$$Re\left[\mu + \sum_{i=1}^{r} \rho_{i} \min_{1 \leq j \leq m_{i}} \frac{d_{j}^{(i)}}{\delta_{j}^{(i)}}\right] > 0$$

$$D_{x}^{\alpha} \Big\{ x^{\mu} (x^{\nu} + \epsilon)^{\lambda} (x^{\nu} + \eta)^{\gamma} \phi_{N}(zx) T_{N'}^{M'_{1}, \cdots, M'_{k}} \begin{pmatrix} y_{1} x^{\lambda_{1}} (x^{\nu} + \epsilon)^{\delta_{1}} (x^{\nu} + \eta)^{\epsilon_{1}} \\ \ddots \\ y_{k} x^{\lambda_{k}} (x^{\nu} + \epsilon)^{\delta_{k}} (x^{\nu} + \eta)^{\epsilon_{k}} \end{pmatrix}$$

$$I\left(\begin{array}{c} z_1 x^{\rho_1} (x^{\upsilon} + \epsilon)^{-\sigma_1} (x^{\upsilon} + \eta)^{-\mu_1} \\ & \ddots \\ & \ddots \\ z_r x^{\rho_r} (x^{\upsilon} + \epsilon)^{-\sigma_r} (x^{\upsilon} + \eta)^{-\mu_r} \end{array}\right)$$

$${}_{P}F_{Q}\left[(A_{P});(B_{Q});-\sum_{j=1}^{t}z_{j}'x^{\delta_{j}'}(x^{\upsilon}+\epsilon)^{-\rho_{j}'}(x^{\upsilon}+\eta)^{-\lambda_{j}'}\right]\right\}$$

$$=\frac{\epsilon^{\lambda}\eta^{\gamma}\prod_{j=1}^{Q}\Gamma(B_{j})}{\prod_{j=1}^{P}\Gamma(A_{j})}\sum_{h=0}^{[N/M]}\sum_{l=0}^{[dh]}\frac{(-dh)_{l}2^{l}c^{-\upsilon'-N+(M-1)h}(\upsilon')_{N+(1-M)h}(a)^{N-Mh}b^{h}(-)^{(d+1)h}2^{l}x^{N-Mh+1}}{l!h!(N-Mh)!}$$

$$\sum_{h_1,\cdots,h_k=0}^{M'\leqslant N'} \left[(-N')_{M'} B(N',h_1,\cdots,h_k) \right] \prod_{j=1}^k \left[\frac{(-N'_j)_{M'_j h_j}}{h_j!} (x^{\lambda_j} \epsilon^{\delta_j} \eta^{\epsilon_j} y_j)^{h_j} \right] \sum_{l_1,l_2=0}^{\infty} \frac{(-x)^{\upsilon l_1+\upsilon l_2}}{\epsilon^{l_1} l_1! \eta^{l_2} l_2!}$$

$$I_{p+P+3,q+Q+3;V}^{0,n+P+3;X} \begin{pmatrix} z_{1}\epsilon^{-\sigma_{1}}\eta^{-\mu_{1}}x^{\rho_{1}} & A, K_{1}:C \\ \vdots & z_{r}\epsilon^{-\sigma_{r}}\eta^{-\mu_{r}}x^{\rho_{r}} \\ z_{1}^{\prime}\epsilon^{-\rho_{1}^{\prime}}\eta^{-\lambda_{1}^{\prime}}x^{\delta_{1}^{\prime}} & z^{\prime}_{1}\epsilon^{-\rho_{1}^{\prime}}\eta^{-\lambda_{1}^{\prime}}x^{\delta_{1}^{\prime}} \\ \vdots & z^{\prime}_{t}\epsilon^{-\rho_{t}^{\prime}}\eta^{-\lambda_{t}^{\prime}}x^{\delta_{t}^{\prime}} & B, K_{2}:D \end{pmatrix}$$
(3.11)

where $M' = \sum_{i=1}^{k} M'_{i} h_{i}$, under the same conditions and notations that (3.8)

Remark: b) If $A_j = B_j = C_j^{(i)} = D_j^{(i)} = 1$, the multivariable I-function defined by Nambisan et al [6] degenere in multivariable H-function defined by Srivastava et al [11], see Jaimini et al [2, page 13].

Concerning the following theorems, the quantities A, B, C and D are similar.

Let
$$A' = (a'_j; \alpha'_j^{(1)}, \cdots, \alpha'_j^{(s)}, 0, \cdots, 0; A'_j)_{1,p'}$$
 (3.12)

$$X' = m'_1, n'_1; \cdots; m'_s, n'_s; V' = p'_1, q'_1; \cdots; p'_s, q'_s$$
(3.13)

$$B' = (b'_j; \beta'_j^{(1)}, \cdots, \beta'_j^{(s)}, 0, \cdots, 0; B'_j)_{1,q'}$$
(3.14)

$$C' = (c'_{j}^{(1)}, \gamma'_{j}^{(1)}; C'_{j}^{(1)})_{1, p'_{1}}; \cdots; (c'_{j}^{(r)}, \gamma^{(s)}_{j}; C'_{j}^{(s)})_{1, p'_{s}}$$
(3.15)

$$D' = (d_{j}^{(1)}, \delta_{j}^{\prime(1)}; D_{j}^{\prime(1)})_{1,q_{1}^{\prime}}; \cdots; (d_{j}^{\prime(s)}, \delta_{j}^{\prime(s)}; D_{j}^{\prime(s)})_{1,q_{s}^{\prime}}$$
(3.16)

$$K_3 = (-\mu; \lambda_1, \cdots, \lambda_s; 1) \text{and} K_4 = (-\mu + \alpha - en' - \epsilon; \lambda_1, \cdots, \lambda_s; 1)$$
(3.17)

$$K_{5} = [1 - A_{j}; 0, \dots, 0, 1, \dots, 1; 1]_{1,P}, (1 + \lambda + \sum_{j=1}^{k} \delta_{j}h_{j}; \sigma_{1}, \dots, \sigma_{r}, \delta_{1}, \dots, \delta_{t}'; 1),$$

$$(1 + \gamma + \sum_{j=1}^{k} \epsilon_{j}h_{j} - l_{2}; \mu_{1}, \dots, \mu_{r}, \epsilon_{1}', \dots, \epsilon_{t}'; 1),$$

$$(Mh - \mu - N - l - \upsilon l_{1} - \upsilon l_{2} - \sum_{j=1}^{k} \eta_{j}h_{j}; \rho_{1}, \dots, \rho_{r}, \rho_{1}', \dots, \rho_{t}'; 1)$$

$$K_{6} = [1 - B_{j}; 0, \dots, 0, 1, \dots, 1; 1]_{1,Q}, (1 + \lambda + \sum_{j=1}^{k} \delta_{j}h_{j}; \sigma_{1}, \dots, \sigma_{r}, \delta_{1}', \dots, \delta_{t}'; 1),$$
(3.18)

$$(1 + \gamma + \sum_{j=1}^{k} \epsilon_{j} h_{j}; \mu_{1}, \cdots, \mu_{r}, \epsilon_{1}', \cdots, \epsilon_{t}'; 1),$$

$$(Mh - N - l - \upsilon l_{1} - \upsilon l_{2} - \sum_{j=1}^{k} \eta_{j} h_{j} + en' + \epsilon; \rho_{1}, \cdots, \rho_{r}, \rho_{1}', \cdots, \rho_{t}'; 1)$$
(3.19)

$$D_x^{\alpha} \Big\{ x^{\mu} (x^{\upsilon} + \epsilon)^{\lambda} (x^{\upsilon} + \eta)^{\gamma} \phi_N(zx) \, S_{N_1', \cdots, N_k'}^{M_1', \cdots, M_k'} \left(\begin{array}{c} y_1 x^{\lambda_1} (x^{\upsilon} + \epsilon)^{\delta_1} (x^{\upsilon} + \eta)^{\epsilon_1} \\ & \ddots \\ & \ddots \\ y_k x^{\lambda_k} (x^{\upsilon} + \epsilon)^{\delta_k} (x^{\upsilon} + \eta)^{\epsilon_k} \end{array} \right)$$

$$I\left(\begin{array}{ccc} z_{1}x^{\rho_{1}}(x^{\upsilon}+\epsilon)^{-\sigma_{1}}(x^{\upsilon}+\eta)^{-\mu_{1}}\\ \vdots\\ z_{r}x^{\rho_{r}}(x^{\upsilon}+\epsilon)^{-\sigma_{r}}(x^{\upsilon}+\eta)^{-\mu_{r}}\end{array}\right)I\left(\begin{array}{ccc} w_{1}x^{\lambda_{1}}\\ \vdots\\ \vdots\\ \vdots\\ w_{s}x^{\lambda_{s}}\end{array}\middle| A^{\prime}:C^{\prime}\\ \vdots\\ \vdots\\ w_{s}x^{\lambda_{s}}}\middle| B^{\prime}:D^{\prime}\right)$$

$${}_{P}F_{Q}\left[(A_{P});(B_{Q});-\sum_{j=1}^{t}z_{j}'x^{\delta_{j}'}(x^{\upsilon}+\epsilon)^{-\rho_{j}'}(x^{\upsilon}+\eta)^{-\lambda_{j}'}\right]\right\} = \frac{\epsilon^{\lambda}\eta^{\gamma}x^{\mu-\alpha}\prod_{j=1}^{Q}\Gamma(B_{j})}{\prod_{j=1}^{P}\Gamma(A_{j})}$$

$$\sum_{h=0}^{[N/M]} \sum_{l=0}^{[dh]} \frac{(-dh)_l x^{N-Mh+l} c^{-\upsilon'-N+(M-1)h} (\upsilon')_{N+(1-M)h} (a)^{N-Mh} b^h (-)^{(d+1)h} 2^l x^{N-Mh+1}}{l!h! (N-Mh)!}$$

$$\sum_{h_1=0}^{[N_1'/M_1']} \cdots \sum_{h_k=0}^{[N_k'/M_k']} A(h_1, \cdots, h_k) \prod_{j=1}^k \left[\frac{(-N_j')_{M_j'h_j}}{h_j!} (x^{\lambda_j} \epsilon^{\delta_j} \eta^{\epsilon_j} y_j)^{h_j} \right] \sum_{n'=-\infty}^{\infty} \sum_{l_1, l_2=0}^{\infty} e\binom{\alpha}{\epsilon n' + e} \frac{(-x)^{\nu l_1 + \nu l_2}}{\epsilon^{l_1} l_1! \eta^{l_2} l_2!}$$

$$I_{p+P+3,q+Q+3;V}^{0,n+P+3;X} \begin{pmatrix} z_{1}\epsilon^{-\sigma_{1}}\eta^{-\mu_{1}}x^{\rho_{1}} \\ \vdots \\ z_{r}\epsilon^{-\sigma_{r}}\eta^{-\mu_{r}}x^{\rho_{r}} \\ z'_{1}\epsilon^{-\rho'_{1}}\eta^{-\lambda'_{1}}x^{\delta'_{1}} \\ \vdots \\ z'_{t}\epsilon^{-\rho'_{t}}\eta^{-\lambda'_{t}}x^{\delta'_{t}} \end{pmatrix} B, K_{6}:D \end{pmatrix} I_{p'+1,q';V'}^{0,n'+1;X'} \begin{pmatrix} w_{1}x^{\lambda_{1}} \\ \vdots \\ \vdots \\ w_{1}x^{\lambda_{1}} \\ \vdots \\ \vdots \\ \vdots \\ w_{2}x^{\lambda_{2}} \\ B',K_{4}:D' \end{pmatrix} (3.20)$$

Provided that

a) $\mu, \eta_i, \delta_i, \epsilon_i, \rho_j, \mu_j, \rho'_l, \delta'_l, \epsilon'_l, \lambda_p > 0, i = 1, \cdots, k; j = 1, \cdots, r; l = 1, \cdots, t; p = 1, \cdots, s$

$$\mathbf{b} \, Re \left[1 + \sum_{i=1}^{r} \rho_{i} \min_{1 \leqslant j \leqslant m_{i}} \frac{d_{j}^{(i)}}{\delta_{j}^{(i)}} \right] > 0 \, ; \, Re \left[\mu + 1 + \sum_{i=1}^{s} \lambda_{i} \min_{1 \leqslant j \leqslant m_{i}'} \frac{d_{j}^{'(i)}}{\delta_{j}^{'(i)}} \right] > 0$$

$$D_x^{\alpha} \Big\{ x^{\mu} (x^{\nu} + \epsilon)^{\lambda} (x^{\nu} + \eta)^{\gamma} \phi_N(zx) T_{N'}^{M'_1, \cdots, M'_k} \begin{pmatrix} y_1 x^{\lambda_1} (x^{\nu} + \epsilon)^{\delta_1} (x^{\nu} + \eta)^{\epsilon_1} \\ & \ddots \\ y_k x^{\lambda_k} (x^{\nu} + \epsilon)^{\delta_k} (x^{\nu} + \eta)^{\epsilon_k} \end{pmatrix}$$

$$I\left(\begin{array}{ccc} z_1 x^{\rho_1} (x^{\upsilon} + \epsilon)^{-\sigma_1} (x^{\upsilon} + \eta)^{-\mu_1} \\ \vdots \\ z_r x^{\rho_r} (x^{\upsilon} + \epsilon)^{-\sigma_r} (x^{\upsilon} + \eta)^{-\mu_r} \end{array}\right) I\left(\begin{array}{ccc} w_1 x^{\lambda_1} \\ \vdots \\ \vdots \\ w_s x^{\lambda_s} \\ w_s x^{\lambda_s} \end{array} \middle| A_1 : C_1 \\ \vdots \\ \vdots \\ B_1 : D_1 \end{array}\right)$$

$${}_{P}F_{Q}\left[(A_{P});(B_{Q});-\sum_{j=1}^{t}z_{j}'x^{\delta_{j}'}(x^{\upsilon}+\epsilon)^{-\rho_{j}'}(x^{\upsilon}+\eta)^{-\lambda_{j}'}\right]\right\} = \frac{\epsilon^{\lambda}\eta^{\gamma}x^{\mu-\alpha}\prod_{j=1}^{Q}\Gamma(B_{j})}{\prod_{j=1}^{P}\Gamma(A_{j})}$$

$$\sum_{h=0}^{\lfloor N/M \rfloor} \sum_{l=0}^{\lfloor dh \rfloor} \frac{(-dh)_l x^{N-Mh+l} c^{-\upsilon'-N+(M-1)h} (\upsilon')_{N+(1-M)h} (a)^{N-Mh} b^h (-)^{(d+1)h} 2^l x^{N-Mh+1}}{l!h! (N-Mh)!}$$

$$\sum_{h_1,\cdots,h_k=0}^{M'\leqslant N'} \left[(-N')_{M'} B(N',h_1,\cdots,h_k) \right] \prod_{j=1}^k \left[\frac{(-N'_j)_{M'_j h_j}}{h_j!} (x^{\lambda_j} \epsilon^{\delta_j} \eta^{\epsilon_j} y_j)^{h_j} \right] \sum_{n'=-\infty}^{\infty} \sum_{l_1,l_2=0}^{\infty} e\binom{\alpha}{\epsilon n'+e} \frac{(-x)^{\nu l_1+\nu l_2}}{\epsilon^{l_1} l_1! \eta^{l_2} l_2!}$$

$$I_{p+P+3,q+Q+3;V}^{0,n+P+3;X}\begin{pmatrix} z_{1}\epsilon^{-\sigma_{1}}\eta^{-\mu_{1}}x^{\rho_{1}} \\ \vdots \\ z_{r}\epsilon^{-\sigma_{r}}\eta^{-\mu_{r}}x^{\rho_{r}} \\ z'_{1}\epsilon^{-\rho'_{1}}\eta^{-\lambda'_{1}}x^{\delta'_{1}} \\ \vdots \\ z'_{t}\epsilon^{-\rho'_{t}}\eta^{-\lambda'_{t}}x^{\delta'_{t}} \\ z'_{t}\epsilon^{-\rho'_{t}}\eta^{-\lambda'_{t}}x^{\delta'_{t}} \end{bmatrix} \mathbf{A}, \mathbf{K}_{5}:C \\ \mathbf{H}_{p'+1,q';V'}^{0,n'+1;X'}\begin{pmatrix} \mathbf{W}_{1}x^{\lambda_{1}} \\ \vdots \\ \mathbf{W}_{s}x^{\lambda_{s}} \\ \mathbf{H}_{1},\mathbf{K}_{3}:C_{1} \\ \vdots \\ \mathbf{W}_{s}x^{\lambda_{s}} \\ \mathbf{H}_{1},\mathbf{K}_{4}:D_{1} \end{pmatrix}$$
(3.21)

where $M' = \sum_{i=1}^{k} M'_{i} h_{i}$, under the same conditions and notations that (3.10) and

Remark concerning the theorems 3 and 4 : If ; $A_j = B_j = C_j^{(i)} = D_j^{(i)} = A'_j = B'_j = C'_j^{(i)} = D'_j^{(i)} = 1$ The multivariable I-function defined by Nambisan et al [6] degenere in the multivariable H-functions defined by Srivastava et al [11] and we obtain the H-function of (r + t) variables and H-function of s variables, see Jaimini et al [2, page 13].

Proofs

To prove the theorem 1, we have

$$\text{L.H.S} = D_x^{\alpha} \left\{ x^{\mu} (x^{\nu} + \epsilon)^{\lambda} (x^{\nu} + \eta)^{\gamma} \phi_N(zx) S_{N_1', \cdots, N_k'}^{M_1', \cdots, M_k'} \begin{pmatrix} y_1 x^{\lambda_1} (x^{\nu} + \epsilon)^{\delta_1} (x^{\nu} + \eta)^{\epsilon_1} \\ & \ddots \\ & \ddots \\ y_k x^{\lambda_k} (x^{\nu} + \epsilon)^{\delta_k} (x^{\nu} + \eta)^{\epsilon_k} \end{pmatrix} \right\}$$

$$I\left(\begin{array}{c} z_1 x^{\rho_1} (x^{\upsilon} + \epsilon)^{-\sigma_1} (x^{\upsilon} + \eta)^{-\mu_1} \\ \vdots \\ z_r x^{\rho_r} (x^{\upsilon} + \epsilon)^{-\sigma_r} (x^{\upsilon} + \eta)^{-\mu_r} \end{array}\right)$$

$${}_{P}F_{Q}\left[(A_{P});(B_{Q});-\sum_{j=1}^{t}z_{j}'x^{\delta_{j}'}(x^{\upsilon}+\epsilon)^{-\rho_{j}'}(x^{\upsilon}+\eta)^{-\lambda_{j}'}\right]\right\}$$
(3.22)

First, using the definitions of general class of polynomials $\phi_n(.)$, $S_{N'_1, \cdots, N'_t}^{M'_1, \cdots, M'_t}[.]$ given in (1.15) and (1.16) respectively and expressing the I-function of r variables and generalized hypergeometric function in Mellin-Barnes contour integral with the help of equations (1.2) and (1.21) respectively, changing the order of summations and integrations (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process), we have :

$$\text{L.H.S} = \frac{\prod_{j=1}^{Q} \Gamma(B_j)}{\prod_{j=1}^{P} \Gamma(A_j)} \sum_{h=0}^{[N/M]} \sum_{l=0}^{[dh]} \frac{(-dh)_l x^{N-Mh+l} c^{-\upsilon'-N+(M-1)h} (\upsilon')_{N+(1-M)h} (a)^{N-Mh} b^h (-)^{(d+1)h} 2^l x^{N-Mh+1}}{l!h! (N-Mh)!}$$

$$\sum_{h_1=0}^{[N_1'/M_1']} \cdots \sum_{h_k=0}^{[N_k'/M_k']} A(h_1, \cdots, h_k) \prod_{j=1}^k \left[\frac{(-N_j')_{M_j'h_j}}{h_j!} y_j^{h_j} \right] \frac{1}{(2\pi\omega)^{r+t}} \int_{L_1} \cdots \int_{L_r} \int_{L_1'} \cdots \int_{L_t'} \xi(S_1, \cdots, S_r)$$

$$\prod_{i=1}^{r} \phi_i(S_i) z_j^{S_j} \frac{\prod_{j=1}^{P} \Gamma(A_j + \sum_{i=1}^{k} s_i)}{\prod_{j=1}^{Q} \Gamma(B_j + \sum_{i=1}^{k} s_i)} \prod_{j=1}^{t} \left[\Gamma(-s_j) (z_j')^{s_j} \right] D_x^{\alpha} \Big\{ x^{\mu + N - Mh + 1 - \sum_{j=1}^{k} \lambda_j h_j + \sum_{j=1}^{r} \rho_j S_j + \sum_{j=1}^{t} \delta_j' s_j + \sum_{j$$

$$(x^{\upsilon}+\epsilon)^{\lambda+\sum_{j=1}^{k}\delta_{j}h_{j}-\sum_{j=1}^{r}\sigma_{j}S_{j}-\sum_{j=1}^{t}\rho_{i}'s_{j}}(x^{\upsilon}+\eta)^{\gamma+\sum_{j=1}^{k}\epsilon_{j}h_{j}-\sum_{j=1}^{r}\mu_{j}S_{j}-\sum_{j=1}^{t}\lambda_{i}'s_{j}}\bigg\}\mathrm{d}S_{1}\cdots\mathrm{d}S_{r}\mathrm{d}s_{1}\cdots\mathrm{d}S_{t}(3.23)$$

Now use the binomial formula : $(x^{\upsilon} + a)^{\lambda} = a^{\lambda} \sum_{m=0}^{\infty} {\lambda \choose m} \left(\frac{x^{\upsilon}}{a}\right)^m; \left|\frac{x^{\upsilon}}{a}\right| < 1$ (3.24)

twice and then the result (1.19) therein. On interpreting the contour integrals into I-function of (r+t) variables with the help of (1.2), we obtain the desired result.

To prove the theorem 2, we use the similar method using the definition of general class of polynomials $T_n^{m_1,\dots,m_k}[.]$.

To prove the theorem 3, we set f(x), g(x) in equation (1.20) as the following

$$f(x) = x^{\mu} I \begin{pmatrix} w_1 x^{\lambda_1} & \mathbf{A} : \mathbf{C} \\ \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ w_s x^{\lambda_s} & \mathbf{B} : \mathbf{D} \end{pmatrix}$$
(3.25)

and
$$g(x) = (x^{\upsilon} + \epsilon)^{\lambda} (x^{\upsilon} + \eta)^{\gamma} \phi_N(zx) S_{N'_1, \cdots, N'_k}^{M'_1, \cdots, M'_k} \begin{pmatrix} y_1 x^{\lambda_1} (x^{\upsilon} + \epsilon)^{\delta_1} (x^{\upsilon} + \eta)^{\epsilon_1} \\ & \ddots \\ & \ddots \\ y_k x^{\lambda_k} (x^{\upsilon} + \epsilon)^{\delta_k} (x^{\upsilon} + \eta)^{\epsilon_k} \end{pmatrix}$$

$$I\begin{pmatrix} z_{1}x^{\rho_{1}}(x^{\upsilon}+\epsilon)^{-\sigma_{1}}(x^{\upsilon}+\eta)^{-\mu_{1}}\\ \vdots\\ z_{r}x^{\rho_{r}}(x^{\upsilon}+\epsilon)^{-\sigma_{r}}(x^{\upsilon}+\eta)^{-\mu_{r}} \end{pmatrix} PF_{Q}\left[(A_{P});(B_{Q});-\sum_{j=1}^{t}z_{j}'x^{\delta_{j}'}(x^{\upsilon}+\epsilon)^{-\rho_{j}'}(x^{\upsilon}+\eta)^{-\lambda_{j}'}\right]$$
(3.26)

We have
$$\text{L.H.S} = \sum_{n'=-\infty}^{\infty} e \binom{\alpha}{en'+\epsilon} D_x^{\alpha-en'-\epsilon} \left[x^{\mu} \aleph(w_1 x^{\lambda_1}, \cdots, w_s x^{\lambda_s}] D_x^{an+\epsilon} \right] \left\{ x^{\mu} (x^{\nu}+\epsilon)^{\lambda} (x^{\nu}+\eta)^{\gamma} \right\}$$

$$\phi_{N}(zx)S_{N_{1}^{\prime},\cdots,N_{k}^{\prime}}^{M_{1}^{\prime},\cdots,M_{k}^{\prime}}\left(\begin{array}{cc}y_{1}x^{\lambda_{1}}(x^{\upsilon}+\epsilon)^{\delta_{1}}(x^{\upsilon}+\eta)^{\epsilon_{1}}\\ & \ddots\\ & \ddots\\ & y_{k}x^{\lambda_{k}}(x^{\upsilon}+\epsilon)^{\delta_{k}}(x^{\upsilon}+\eta)^{\epsilon_{k}}\end{array}\right)I\left(\begin{array}{cc}z_{1}x^{\rho_{1}}(x^{\upsilon}+\epsilon)^{-\sigma_{1}}(x^{\upsilon}+\eta)^{-\mu_{1}}\\ & \ddots\\ & \ddots\\ & z_{r}x^{\rho_{r}}(x^{\upsilon}+\epsilon)^{-\sigma_{r}}(x^{\upsilon}+\eta)^{-\mu_{r}}\end{array}\right)$$

$${}_{P}F_{Q}\left[(A_{P});(B_{Q});-\sum_{j=1}^{t}z_{j}'x^{\delta_{j}'}(x^{\upsilon}+\epsilon)^{-\rho_{j}'}(x^{\upsilon}+\eta)^{-\lambda_{j}'}\right]\right\}$$
(3.27)

Now use the Lemme and the theorem 1, we obtain the theorem 3

To prove the theorem 4, we use the similar method using the definition of general class of polynomials $T_n^{m_1, \dots, m_k}[.]$.

4.Conclusion

In this paper we have evaluated a generalized contour integral involving the multivariable I-functions defined by Nambisan et al [6], two classes of polynomials and the generalized hypergeometric function. The four formulaes established in this paper is very general nature. Thus, the results established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

[1] B. L. J. Braaksma, "Asymptotic expansions and analytic continuations for a class of Barnes integrals," Compositio Mathematical, vol. 15, pp. 239–341, 1964.

[2] Jaimini B.B. and Gautam S. Fractional derivatives in general classes of polynomials and special functions representable by Mellin-Barnes type contour integrals. Jnanabha Vol 36 (2006), page 9-20.

[3] Oldham K.B.and Spanier J. The fractional calculus. Academic Press, New York 1974.

[4] Osler T.J. Fractional derivatives and Leibnitz rule. Amer. Math. Montly. Vol 78 (1971) ,page 645-649.

[5] Pathan M.A. And Khan M.A. On polynomials associated with Humbert's polynomials . Publ. De. L'institut. Mathématiques. Nouvelles. Série, 62(76) 1997, page 53-62.

[6] Prathima J. Nambisan V. and Kurumujji S.K. A Study of I-function of Several Complex Variables, International Journalof Engineering Mathematics Vol(2014), 2014 page 1-12

[7] Samko S.G. Kilbas A.A. and Marichev O.I. Fractional integral and derivatives : Theory and applications, Gordon and Breach Sciences Publishers, Reading. Tokyo, Pais, Berlin and Langhorne (Pennsylvania), 1993

[8] Srivastava H.M. A multilinear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial, Pacific. J. Math. 177(1985), page183-191

[9] Srivastava H.M and Garg M. Some integrals involving a general class of polynomials and the multivariable H-function. Rev. Roumaine. Phys.32 1987 ,page 685-692.

[10] Srivastava H.M. and Karlsson P.W. Multiple Gaussian Hypergeometric series. Ellis.Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.

[11] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.

Personal adress : 411 Avenue Joseph Raynaud Le parc Fleuri , Bat B 83140 , Six-Fours les plages Tel : 06-83-12-49-68 Department : VAR Country : FRANCE